


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



26

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C07D 471/16, A61K 31/395, A61P 43/00, C07D 498/16, 513/16 // (C07D 471/16, 223:00, 221:00, 209:00) (C07D 471/16, 223:00, 209:00, 209:00) (C07D 498/16, 265:00, 223:00, 209:00) (C07D 513/16, 279:00, 223:00, 209:00)</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 00/64899</b>  (43) International Publication Date: 2 November 2000 (02.11.00)</p>
<p>(21) International Application Number: PCT/US00/10600  (22) International Filing Date: 20 April 2000 (20.04.00)  (30) Priority Data: 60/130,811 23 April 1999 (23.04.99) US  (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/130,811 (CIP) Filed on 23 April 1999 (23.04.99)  (71) Applicant (for all designated States except US): PHARMACIA &amp; UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p>		<p>(71)(72) Applicants and Inventors: HESTER, Jackson, B., Jr. [US/US]; 9219 East ML Avenue, Galesburg, MI 49053 (US). ROGERS, Bruce, N. [US/US]; 5860 Tradewind Drive, Portage, MI 49024 (US). JACOBSEN, E., Jon [US/US]; 74 South Lake Doster Drive, Plainwell, MI 49080 (US). ENNIS, Michael, Dalton [US/US]; 2416 Deep Forest Court, Portage, MI 49002 (US). ACKER, Brad, A. [US/US]; 1734 Tanager Lane, Kalamazoo, MI 49009 (US). VANDER VELDE, Susan, L. [US/US]; 6125 Fairgrove, Kalamazoo, MI 49009 (US).  (74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg, Woessner &amp; Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).  (81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: TETRACYCLIC AZEPINOINDOLE COMPOUNDS AS 5-HT RECEPTOR LIGANDS  (57) Abstract  The present invention provides compounds of formula (I), wherein R<sub>2</sub> is hydrogen, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkanoyl, arylcarbonyl, aryl, (aryl)C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxycarbonyl, aryloxycarbonyl, arylsulfonyl, or (aryl)C<sub>1-7</sub>alkoxycarbonyl; X and Y together are 2, 3 or 4 membered saturated or partially unsaturated chain. These compounds are 5-HT ligands, and are useful for treating diseases wherein modulation of 5-HT activity is desired.</p> <div style="text-align: center;">  <p>(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## TETRACYCLIC AZEPINOINDOLE COMPOUNDS AS 5-HT RECEPTOR LIGANDS

5

## PRIORITY OF INVENTION

This application claims priority from United States Provisional Application Number 60/130,881, filed 23 April 1999.

## FIELD OF THE INVENTION

The present invention provides tetracyclic 1,2,3,4,5,6-  
10 hexahydroazepino-[4,5-b]indole derivatives having a ring connecting position 6 (N-6) and position 7 (C-7), and more specifically, provides compounds of formula (I) described hereinbelow. These compounds are 5-HT ligands, and are useful for treating diseases wherein modulation of 5-HT activity is desired.

## BACKGROUND OF THE INVENTION

15

Serotonin has been implicated in a number of diseases and conditions which originate in the central nervous system. These include diseases and conditions related to sleeping, eating, perceiving pain, controlling body temperature, controlling blood pressure, depression, anxiety, schizophrenia, and other bodily states. R. W. Fuller, *Biology of Serotonergic Transmission*, 221  
20 (1982); D. J. Boullin, *Serotonin in Mental Abnormalities* 1:316 (1978); J. Barchas, et al., *Serotonin and Behavior*, (1973). Serotonin also plays an important role in peripheral systems, such as the gastrointestinal system, where it has been found to mediate a variety of contractile, secretory, and electrophysiologic effects.

25

As a result of the broad distribution of serotonin within the body, there is a tremendous interest in drugs that affect serotonergic systems. In particular, receptor-specific agonists and antagonists are of interest for the treatment of a wide range of disorders, including anxiety, depression, hypertension, migraine, obesity, compulsive disorders, schizophrenia, autism,  
30 neurodegenerative disorders (e.g. Alzheimer's disease, Parkinsonism, and

Huntington's chorea), and chemotherapy-induced vomiting. M. D. Gershon, et al., *The Peripheral Actions of 5-Hydroxytryptamine*, 246 (1989); P. R. Saxena, et al., *Journal of Cardiovascular Pharmacology*, 15:Supplement 7 (1990).

The major classes of serotonin receptors (5-HT<sub>1-7</sub>) contain  
5 fourteen to eighteen separate receptors that have been formally classified. See  
Glennon, et al., *Neuroscience and Behavioral Reviews*, 1990, 14, 35; and D.  
Hoyer, et al. *Pharmacol. Rev.* 1994, 46, 157-203. Recently discovered  
information regarding subtype identity, distribution, structure, and function  
suggests that it is possible to identify novel, subtype specific agents, having  
10 improved therapeutic profiles (e.g. fewer side effects).

For example, the 5-HT<sub>2</sub> family of receptors is comprised of 5-  
HT<sub>2A</sub>, 5-HT<sub>2B</sub>, and 5-HT<sub>2C</sub> subtypes, which have been grouped together on the  
basis of primary structure, secondary messenger system, and operational profile.  
All three subtypes are G-protein coupled, activate phospholipase C as a  
15 principal transduction mechanism, and contain a seven-transmembrane domain  
structure. There are distinct differences in the distribution of the three 5-HT<sub>2</sub>  
subtypes. The 5-HT<sub>2B</sub> and 5-HT<sub>2A</sub> receptors are widely distributed in the  
periphery, while the 5-HT<sub>2C</sub> receptor has been found only in the central nervous  
system, being highly expressed in many regions of the human brain. See G.  
20 Baxter, et al. *Trends in Pharmacol. Sci.* 1995, 16, 105-110.

Subtype 5-HT<sub>2A</sub> has been associated with effects including  
vasoconstriction, platelet aggregation, and bronchoconstriction, while subtype 5-  
HT<sub>2C</sub> has been associated with diseases that include depression, anxiety,  
obsessive compulsive disorder, panic disorders, phobias, psychiatric syndromes,  
25 and obesity. Very little is known about the pharmacologic role of the 5-HT<sub>2B</sub>  
receptor. See F. Jenck, et al., *Exp. Opin. Invest. Drugs*, 1998, 7, 1587-1599; M.  
Bos, et al., *J. Med. Chem.*, 1997, 40, 2762-2769; J.R. Martin, et al., *The Journal  
of Pharmacology and Experimental Therapeutics*, 1998, 286, 913-924; S.M.  
Bromidge, et al., *J. Med. Chem.*, 1998, 41, 1598-1612; G.A. Kennett, *Drugs*,

1998, 1, 4, 456-470; and A. Dekeyne, et al., *Neuropharmacology*, 1999, 38, 415-423.


United States Patent Number 3,676,558, issued July 11, 1972, discloses compositions comprising specific 6-alkyl-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole compounds. The compositions are reported to be useful to suppress hunger in mammals. This patent also discloses a method for inducing anorexia in obese subjects to produce weight loss. The azepino[4,5-b]indole compounds disclosed in this patent lack the ring connecting the 6-position and the 7-position that is present in the compounds of the instant invention.

United States Patent Number 3,839,357, issued October 1, 1974, discloses specific 1,2,3,4,5,6-hexahydroazepino[4,5-b]indole compounds, which are reported to have sedative or tranquilizing action. The azepino[4,5-b]indole compounds disclosed in this patent also lack the ring connecting the 6-position and the 7-position that is present in the compounds of the instant invention.

There is currently a need for pharmaceutical agents that are useful to treat diseases and conditions that are associated with 5-HT receptors.

#### SUMMARY OF THE INVENTION

In accordance with the present invention, novel compounds which demonstrate useful biological activity, and particularly activity as 5-HT receptor ligands, are provided. Thus, the present invention provides a compound of formula I:



(I)

wherein,

- each  $R_1$  is independently hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl,  $-S(O)_mNR_4R_b$ ,  $NR_4R_b$ ,  $-S(O)_mR_c$ , or  $-C(=O)NR_4R_b$ , wherein any  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, or  $C_{1-7}$ alkanoyloxy of  $R_1$  is optionally partially unsaturated and is optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy,  $-S(O)_mR_c$ ,  $-S(O)_mNR_4R_b$ ,  $NR_4R_b$ , or  $-C(=O)NR_4R_b$ ;
- $R_2$  is hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl, arylcarbonyl, aryl, (aryl) $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxycarbonyl, aryloxycarbonyl, arylsulfonyl, or (aryl) $C_{1-7}$ alkoxycarbonyl;
- X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl ( $S(O)_2$ -), or  $NR_4$  in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), thioxo (=S),  $-NR_4R_b$ ,  $-S(O)_pR_c$ , or  $-OR_p$ , or with one or two substituents independently selected from the group consisting of  $C_{1-7}$ alkyl, ( $C_{1-7}$ alkoxy) $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, and (aryl)oxy $C_{1-7}$ alkyl; or wherein the chain is optionally substituted on a carbon with a 4, 5, or 6 membered spirocyclic carbon ring; or wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain (e.g.  $-CH_2CH_2-$ ,  $-CH_2CH_2CH_2-$ , or  $-CH_2CH_2CH_2CH_2-$ ) forming a ring that is fused to the ring comprising X and Y;
- each m is independently 0, 1, or 2;
- n is 0, 1, 2, or 3;
- p is 0, 1, or 2;
- each  $R_4$  and  $R_b$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl; or  $R_4$  and  $R_b$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

each  $R_c$  and  $R_d$  is independently hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, arylcarbonyl, heteroarylcarbonyl, aryloxycarbonyl, or heteroaryloxycarbonyl; or  $R_c$  and  $R_d$  together with the nitrogen to which they are attached form a  
5 pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

each  $R_e$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl;

$R_f$  is hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, or is a bond to a 2, 3, or 4 membered alkylene chain that  
10 forms a ring that is fused to the ring comprising X and Y;

each  $R_g$  and  $R_h$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl; or  $R_g$  and  $R_h$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

15  $R_i$  is  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl; and

$R_j$  is hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl;

wherein any aryl or heteroaryl ring of  $R_1$ ,  $R_2$ , X, Y,  $R_a$ - $R_b$ , or  $R_g$ - $R_h$   
20 is optionally substituted with one or more (e.g. 1, 2, 3, or 4) substituents independently selected from halo, hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy, phenyl, sulfonyl,  $NR_jR_k$ , or - $C(=O)NR_jR_k$ ; wherein each  $R_j$  and  $R_k$  is independently hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, aryl, (aryl) $C_{1-7}$ alkyl, arylcarbonyl, or  
25 aryloxycarbonyl; or  $R_j$  and  $R_k$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

or a pharmaceutically acceptable salt thereof;

provided Y is not oxy, thio, sulfinyl, or  $NR_i$ ; and

provided X and Y together are not a 2-membered unsaturated  
30 chain; and

provided no carbon of X and Y is bonded to more than one oxy,  
thio, sulfinyl, or NR<sub>r</sub>.

In another aspect, the present invention also provides:

a pharmaceutical composition comprising a compound of formula

5 I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically  
acceptable excipient (the composition preferably comprises a therapeutically  
effective amount of the compound or salt),

a method for treating a disease or condition in a mammal (e.g. a  
human) wherein a 5-HT receptor is implicated and modulation of a 5-HT  
10 function is desired comprising administering a therapeutically effective amount  
of a compound of formula I, or a pharmaceutically acceptable salt thereof to the  
mammal,

a method for treating or preventing a disease or disorder of the  
central nervous system in a mammal comprising administering a therapeutically  
15 effective amount of a compound of formula I, or a pharmaceutically acceptable  
salt thereof to the mammal,

a compound of formula (I) or a pharmaceutically acceptable salt  
thereof for use in medical diagnosis or therapy (e.g. the treatment or prevention  
of 5-HT related disease such as anxiety, obesity, depression, or a stress related  
20 disease),

the use of a compound of formula I, or a pharmaceutically  
acceptable salt thereof to prepare a medicament useful for treating or preventing  
a disease or disorder of the central nervous system in a mammal, and

a method for modulating 5-HT receptor function, comprising  
25 administering an effective modulatory amount of a compound of formula I, or a  
pharmaceutically acceptable salt thereof.

The invention also provides novel intermediates and processes  
disclosed herein that are useful for preparing compounds of formula I.



## DESCRIPTION OF THE FIGURES

Figures 1-6 illustrate synthetic processes and intermediates useful for preparing compounds of the invention.

## DETAILED DESCRIPTION OF THE INVENTION

5           The compounds of the invention are useful for treating or preventing diseases or disorders of the central nervous system. Specific diseases or disorders of the central nervous system for which a compound of formula I may have activity include, but are not limited to: obesity, depression, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional  
10   disorder, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, a stress induced problem with the urinary, gastrointestinal or cardiovascular system (e.g., stress incontinence), neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine, headaches,  
15   cluster headaches, sexual dysfunction in a mammal (e.g. a human), addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to general medical conditions, attention-deficit hyperactivity disorder, behavioral disturbance (including agitation in conditions associated  
20   with diminished cognition (e.g., dementia, mental retardation or delirium)), bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, movement disorder (e.g., Huntington's disease or Tardive Dyskinesia),  
25   oppositional defiant disorder, peripheral neuropathy, post-traumatic stress disorder, premenstrual dysphoric disorder, a psychotic disorder (brief and long duration disorders, psychotic disorder due to medical condition, psychotic disorder NOS), mood disorder (major depressive or bipolar disorder with psychotic features) seasonal affective disorder, a sleep disorder, a specific

development disorder, agitation disorder, selective serotonin reuptake inhibition (SSRI) "poop out" syndrome or a Tic disorder (e.g., Tourette's syndrome).

The following definitions are used, unless otherwise described:

halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, etc. denote both straight  
5 and branched groups; but reference to an individual radical such as "propyl"  
embraces only the straight chain radical, a branched chain isomer such as  
"isopropyl" being specifically referred to. When alkyl can be partially  
unsaturated, the alkyl chain may comprise one or more (e.g. 1, 2, 3, or 4) double  
or triple bonds in the chain.

10 Aryl denotes a phenyl radical or an ortho-fused bicyclic  
carbocyclic radical having about nine to ten ring atoms in which at least one ring  
is aromatic. Heteroaryl denotes a radical of a monocyclic aromatic ring  
containing five or six ring atoms consisting of carbon and 1, 2, 3, or 4  
heteroatoms each selected from the group consisting of non-peroxide oxygen,  
15 sulfur, and N(X) wherein X is absent or is H, O, C<sub>1-4</sub>alkyl, phenyl or benzyl, as  
well as a radical of an ortho-fused bicyclic heterocycle of about eight to ten ring  
atoms derived therefrom, particularly a benz-derivative or one derived by fusing  
a propylene, trimethylene, or tetramethylene diradical thereto.

It will be appreciated by those skilled in the art that compounds of  
20 the invention having a chiral center may exist in and be isolated in optically  
active and racemic forms. Some compounds may exhibit polymorphism. It is to  
be understood that the present invention encompasses any racemic, optically-  
active, polymorphic, tautomeric, or stereoisomeric form, or mixture thereof, of a  
compound of the invention, which possesses the useful properties described  
25 herein, it being well known in the art how to prepare optically active forms (for  
example, by resolution of the racemic form by recrystallization techniques, by  
synthesis from optically-active starting materials, by chiral synthesis, or by  
chromatographic separation using a chiral stationary phase) and how to  
determine 5-HT activity using the standard tests which are well known in the art.

The carbon atom content of various hydrocarbon-containing moieties is indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix C<sub>i-j</sub> indicates a moiety of the integer "i" to the integer "j" carbon atoms, inclusive. Thus, for example, C<sub>1-7</sub>alkyl refers to alkyl of one to seven carbon atoms, inclusive.

The compounds of the present invention are generally named according to the IUPAC or CAS nomenclature system. Abbreviations which are well known to one of ordinary skill in the art may be used (e.g. "Ph" for phenyl, "Me" for methyl, "Et" for ethyl, "h" for hour or hours and "rt" for room temperature).

Specific and preferred values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents

Specifically, C<sub>1-7</sub>alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, hexyl, or heptyl; C<sub>1-7</sub>alkoxy can be methoxy, ethoxy, propoxy, isopropoxy, butoxy, iso-butoxy, sec-butoxy, pentoxy, 3-pentoxy, hexyloxy, 1-methylhexyloxy, or heptyloxy; C<sub>1-7</sub>alkanoyl can be acetyl, propanoyl, butanoyl, pentanoyl, 4-methylpentanoyl, hexanoyl, or heptanoyl; C<sub>1-7</sub>alkoxycarbonyl can be methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, hexyloxycarbonyl, or heptyloxycarbonyl; C<sub>1-7</sub>alkanoyloxy can be acetoxo, propanoyloxy, butanoyloxy, isobutanoyloxy, pentanoyloxy, hexanoyloxy, or heptanoyloxy; aryl can be phenyl, indenyl, or naphthyl; and heteroaryl can be furyl, imidazolyl, triazolyl, triazinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, pyrrolyl, pyrazinyl, tetrazolyl, pyridyl, (or its N-oxide), thienyl, pyrimidinyl (or its N-oxide), indolyl, isoquinolyl (or its N-oxide) or quinolyl (or its N-oxide).

A specific value for R<sub>1</sub> is hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl, C<sub>1-</sub>

- ,alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, aryl, heteroaryl, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>, NR<sub>c</sub>R<sub>d</sub>,  
 -S(O)<sub>m</sub>R<sub>e</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>, wherein any C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl,  
 C<sub>1-7</sub>alkoxycarbonyl, or C<sub>1-7</sub>alkanoyloxy of R<sub>1</sub> is optionally substituted with aryl,  
 aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano, C<sub>1-7</sub>alkoxy,  
 5 C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, -S(O)<sub>m</sub>R<sub>e</sub>, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>,  
 NR<sub>c</sub>R<sub>d</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>.

- A specific value for R<sub>1</sub> is hydroxy, nitro, halo, cyano,  
 trifluoromethyl, trifluoromethoxy, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>  
 ,alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, aryl, heteroaryl, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>, NR<sub>c</sub>R<sub>d</sub>,  
 10 -S(O)<sub>m</sub>R<sub>e</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>, wherein any C<sub>1-7</sub>alkyl or C<sub>1-7</sub>alkoxy of R<sub>1</sub> is  
 optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy,  
 nitro, halo, cyano, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy,  
 -S(O)<sub>m</sub>R<sub>e</sub>, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>, NR<sub>c</sub>R<sub>d</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>.

- A specific value for R<sub>1</sub> is hydroxy, nitro, halo, cyano,  
 15 trifluoromethyl, trifluoromethoxy, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>  
 ,alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, aryl, heteroaryl, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>, NR<sub>c</sub>R<sub>d</sub>,  
 -S(O)<sub>m</sub>R<sub>e</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>, wherein any C<sub>1-7</sub>alkyl is optionally substituted with  
 aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano, C<sub>1-7</sub>alkoxy,  
 C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, -S(O)<sub>m</sub>R<sub>e</sub>, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>,  
 20 NR<sub>c</sub>R<sub>d</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>.

- A specific value for R<sub>1</sub> is hydroxy, nitro, halo, cyano,  
 trifluoromethyl, trifluoromethoxy, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy, C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>  
 ,alkoxycarbonyl, C<sub>1-7</sub>alkanoyloxy, aryl, heteroaryl, -S(O)<sub>m</sub>NR<sub>a</sub>R<sub>b</sub>, NR<sub>c</sub>R<sub>d</sub>,  
 -S(O)<sub>m</sub>R<sub>e</sub>, or -C(=O)NR<sub>a</sub>R<sub>b</sub>.

- 25 A specific value for R<sub>1</sub> is independently C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkoxy,  
 trifluoromethyl, or halo.

A specific value for R<sub>2</sub> is hydrogen.

A specific value for R<sub>2</sub> is C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, arylcarbonyl,  
 (aryl)C<sub>1-2</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, aryloxycarbonyl, arylsulfonyl, or

(aryl)methoxycarbonyl, wherein any aryl is optionally substituted with 1, 2, or 3 substituents independently selected from C<sub>1-4</sub>alkyl and trifluoromethyl.

A specific value for R<sub>2</sub> is methyl, ethyl, propyl, isopropyl, acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, benzyl, or *p*-toluenesulfonyl.

5 A specific value for n is 1, 2, or 3.

A specific value for n is 0.

A specific group of compounds are compounds of formula (I) wherein n is 0. It will be clear to one skilled in the art that when n is 0, the benz ring of the indole in formula (I) is substituted with hydrogens.

10 A specific group of compounds are compounds of formula I wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo  
15 (=O), thioxo (=S), -NR<sub>q</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, or -OR<sub>v</sub>, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.

A specific group of compounds are compounds of formula I  
20 wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo  
(=O), thioxo (=S), -NR<sub>q</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, or -OR<sub>v</sub>.

25 A specific group of compounds are compounds of formula I wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with one or  
30 two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl,

(C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.

A specific group of compounds are compounds of formula I wherein X and Y together are a 2, 3, or 4 membered chain comprising one or  
5 more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), hydroxy, or C<sub>1-7</sub>alkoxy, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl and  
10 (aryl)oxyC<sub>1-7</sub>alkyl; and wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain forming a ring that is fused to the ring comprising X and Y.

A specific group of compounds are compounds of formula I wherein X and Y together are a 2, 3, or 4 membered chain comprising one or  
15 more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), hydroxy, (aryl)oxy, heteroaryl(oxy), or C<sub>1-7</sub>alkoxy, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl,  
20 heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl; and wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain forming a ring that is fused to the ring comprising X and Y.

A specific group of compounds are compounds of formula I wherein X and Y together are a 2, 3, or 4 membered carbon chain wherein the  
25 chain is optionally substituted on each carbon with oxo (=O), hydroxy, or C<sub>1-7</sub>alkoxy, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl and (aryl)oxyC<sub>1-7</sub>alkyl.

A specific group of compounds are compounds of formula I  
30 wherein X and Y together are a 2, 3, or 4 membered carbon chain wherein the

chain is optionally substituted on each carbon with oxo (=O), hydroxy, (aryl)oxy, heteroaryl(oxy) or C<sub>1-7</sub>alkoxy, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.

5 A specific group of compounds are compounds of formula I wherein X and Y together are a 2 or 3 membered carbon chain optionally substituted on each carbon with oxo or hydroxy, or with one or two C<sub>1-7</sub>alkyl.

A specific group of compounds are compounds of formula I wherein X is -O-, -S-, or -C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein R<sub>g</sub> and R<sub>h</sub> are each independently  
10 hydrogen, C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl or (aryl)oxyC<sub>1-7</sub>alkyl, or R<sub>g</sub> and R<sub>h</sub> together are oxo.

A specific group of compounds are compounds of formula I wherein Y is -C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)-, or  
15 -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, and each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

A specific group of compounds are compounds wherein X is -O-, -S-, or -C(R<sub>g</sub>)(R<sub>h</sub>)-; and Y is -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, or -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

20 A specific group of compounds are compounds wherein X is -O- or -S-; and Y is -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)-, or -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

A specific group of compounds are compounds wherein X and Y together are -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -  
25 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>g</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
30 -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>g</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,

-NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)OC(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)OC(=O)-, -OCH(R<sub>g</sub>)C(=O)-, or -OCH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-; wherein each R<sub>g</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl or (aryl)oxyC<sub>1-7</sub>alkyl.

- 5 A specific group of compounds are compounds wherein X and Y together are -CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH=CHCH<sub>2</sub>-, -CH=CHCH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH=CHCH<sub>2</sub>-, -O-CH<sub>2</sub>CH<sub>2</sub>-, -O-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -S-CH<sub>2</sub>CH<sub>2</sub>-, -S-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -S(O)-CH<sub>2</sub>CH<sub>2</sub>-, -S(O)-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -S(O)<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>-, -S(O)<sub>2</sub>-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-,
- 10 -NR<sub>r</sub>-CH<sub>2</sub>CH<sub>2</sub>-, -NR<sub>r</sub>-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>C(=O)-, -CH(R<sub>g</sub>)CH<sub>2</sub>C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>C(=O)-, -CH<sub>2</sub>OC(=O)-, -CH(R<sub>g</sub>)CH<sub>2</sub>OC(=O)-, -OCH<sub>2</sub>C(=O)-, or -OCH<sub>2</sub>CH<sub>2</sub>C(=O)-; wherein each R<sub>g</sub> is independently -NR<sub>r</sub>R<sub>r</sub>-, -S(O)<sub>p</sub>R<sub>r</sub>-, or -OR<sub>r</sub>-.

- A specific group of compounds are compounds wherein X and Y
- 15 together are -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)OC(=O)-, -OCH(R<sub>g</sub>)C(=O)-; wherein each R<sub>g</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

- 20 A specific group of compounds are compounds wherein X and Y together are -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, or -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-; wherein each R<sub>g</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, or together with an R<sub>g</sub> on
- 25 an adjacent carbon atom forms a fused 4, 5, or 6, membered carbocyclic ring.

A specific group of compounds are compounds wherein X and Y together are -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-; wherein each R<sub>g</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl, or (aryl)C<sub>1-7</sub>alkyl.



A specific group of compounds are compounds wherein X and Y together are  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)\text{H}-$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)\text{HCH}_2-$ ,  $-\text{C}(\text{CH}_3)\text{HCH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)\text{H}-$ ,  $-\text{C}(\text{CH}_3)\text{HC}(\text{CH}_3)\text{H}-$ ,  $-\text{CH}(\text{R}_p)\text{CH}(\text{R}_p)-$ ,  $-\text{O}-\text{CH}_2\text{CH}_2-$ ,  $-\text{O}-\text{C}(\text{CH}_3)\text{HCH}_2-$ , or  $-\text{S}-\text{CH}_2\text{CH}_2-$ .

- 5 A specific group of compounds are compounds wherein X and Y together are  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}=\text{CHCH}_2-$ ,  $-\text{O}-\text{CH}_2\text{CH}_2-$ ,  $-\text{S}-\text{CH}_2\text{CH}_2-$ ,  $-\text{S}(\text{O})-\text{CH}_2\text{CH}_2-$ ,  $-\text{S}(\text{O})_2-\text{CH}_2\text{CH}_2-$ ,  $-\text{NR}_r\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(=\text{O})-$ ,  $-\text{CH}_2\text{CH}_2\text{C}(=\text{O})-$ ,  $-\text{CH}_2\text{OC}(=\text{O})-$ , or  $-\text{OCH}_2\text{C}(=\text{O})-$ .

- A specific group of compounds are compounds of formula (I)  
 10 wherein X and Y together are  $-\text{CH}(\text{R}_p)\text{CH}_2-$ ,  $-\text{CH}(\text{R}_p)\text{CH}(\text{R}_p)\text{CH}_2-$ , or  $-\text{O}-\text{CH}_2\text{CH}_2-$ , wherein each  $\text{R}_p$  is independently  $-\text{NR}_q\text{R}_r$ ,  $-\text{S}(\text{O})_p\text{R}_s$ ,  $-\text{OR}_v$ ,  $\text{C}_{1-7}\text{alkyl}$ ,  $(\text{C}_{1-7}\text{alkoxy})\text{C}_{1-7}\text{alkyl}$ , aryl, (aryl) $\text{C}_{1-7}\text{alkyl}$ , heteroaryl, (heteroaryl) $\text{C}_{1-7}\text{alkyl}$ , or (aryl)oxy $\text{C}_{1-7}\text{alkyl}$ .

- A specific group of compounds are compounds of formula (I)  
 15 wherein X and Y together are  $-\text{C}(=\text{O})\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(=\text{O})-$ ,  $-\text{C}(=\text{S})\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(=\text{S})-$ ,  $-\text{C}(=\text{O})\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(=\text{O})\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{C}(=\text{O})-$ ,  $-\text{C}(=\text{S})\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(=\text{S})\text{CH}_2-$ , or  $-\text{CH}_2\text{CH}_2\text{C}(=\text{S})-$ .

A specific group of compounds are compounds wherein n is 1 and  $\text{R}_1$  is  $\text{C}_{1-7}\text{alkyl}$ ,  $\text{C}_{1-7}\text{alkoxy}$ , or halo.

- 20 A specific group of compounds are compounds wherein n is 1 and  $\text{R}_1$  is methyl, methoxy, chloro, or fluoro.

- A specific group of compounds are compounds wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy ( $-\text{O}-$ ),  
 25 thio ( $-\text{S}-$ ), sulfinyl ( $-\text{SO}-$ ), sulfonyl ( $\text{S}(\text{O})_2-$ ), or  $\text{NR}_t$  in the chain; wherein the chain is optionally substituted on each carbon with oxo ( $=\text{O}$ ), hydroxy, or  $\text{C}_{1-7}\text{alkoxy}$ , or with one or two substituents independently selected from the group consisting of  $\text{C}_{1-7}\text{alkyl}$ ,  $(\text{C}_{1-7}\text{alkoxy})\text{C}_{1-7}\text{alkyl}$ , aryl, (aryl) $\text{C}_{1-7}\text{alkyl}$ , heteroaryl, (heteroaryl) $\text{C}_{1-7}\text{alkyl}$ , and (aryl)oxy $\text{C}_{1-7}\text{alkyl}$ ; or wherein the chain is  
 30 optionally substituted on a carbon with a 4, 5, or 6 membered spirocyclic carbon

ring; or wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain (e.g.  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ , or  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ) forming a ring that is fused to the ring comprising X and Y;

When X and Y together, or  $\text{R}_1$ , is a "partially unsaturated" group, such group may comprise one or more (e.g. 1 or 2) carbon-carbon double or triple bonds. For example, when  $\text{R}_1$  is a partially unsaturated  $\text{C}_{1-7}$ alkyl, it can be vinyl, allyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1,3-butadienyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2,4-hexadienyl, 5-hexenyl, ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 5-hexene-1-ynyl, 2-hexynyl, 3-hexynyl, 3-hexen-5-ynyl, 4-hexynyl, or 5-hexynyl.

Specifically, the invention also provides a method for treating or preventing anxiety, obesity, depression, schizophrenia, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, a stress induced problem with the gastrointestinal or cardiovascular system, or sexual dysfunction in a mammal (e.g. a human) comprising administering a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof to the mammal.

Specifically, the invention also provides a method of treating or preventing anxiety, obesity, depression, or a stress related disease, comprising administering to a mammal (e.g. a human) in need of such treatment, a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

Specifically, the invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof to prepare a medicament for treating or preventing anxiety, obesity, depression, schizophrenia, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress

syndrome, immune system depression, a stress induced problem with the gastrointestinal or cardiovascular system, or sexual dysfunction in a mammal (e.g. a human),.

Specifically, the invention also provides the use of a compound of  
5 formula (I) or a pharmaceutically acceptable salt thereof to prepare a medicament for treating or preventing anxiety, obesity, depression, or a stress related disease in a mammal (e.g. a human).

The invention also provides processes and intermediates useful for preparing compounds of formula (I). For example, an intermediate useful  
10 for preparing a compound of formula (I) wherein  $R_2$  is hydrogen, is a corresponding compound of formula (I) wherein  $R_2$  is a suitable protecting group. Thus the invention provides a compound of formula (I) wherein  $R_2$  is a suitable protecting group, and wherein  $R_1$ , X, Y, and n have any of the values, specific values or preferred values defined herein. Suitable amine protecting  
15 groups, as well as methods for their preparation and removal are well known in the art, for example, see Greene, T.W.; Wutz, P.G.M. "Protecting Groups In Organic Synthesis" third edition, 1999, New York, John Wiley & sons, Inc. Preferred protecting groups include benzyloxycarbonyl (CBZ) and benzoyl.

The invention also provides intermediate compounds of formula  
20 3, 9, 10, 11, 13, 15, and 17-20 as shown in Figures 1-6, wherein  $R_2$  is a protecting group.

The invention also provides intermediate salts that are useful for preparing or purifying compounds of formula (I). Suitable methods for preparing salts are known in the art and are disclosed herein. For example, the  
25 preparation of an oxylate salt is shown in Example 41. As will be apparent to one skilled in the art, such salts can be converted to the corresponding free-base or to another salt using known methods.

For example, compounds of formula I wherein  $R_2$  is  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl, arylcarbonyl, (aryl) $C_{1-2}$ alkyl,  $C_{1-4}$ alkoxycarbonyl, aryloxycarbonyl,  
30 arylsulfonyl, or (aryl)methoxycarbonyl, wherein any aryl is optionally substituted

with 1, 2, or 3 substituents independently selected from C<sub>1-4</sub>alkyl and trifluoromethyl, are particularly useful as intermediates for preparing corresponding compounds of formula I wherein R<sub>2</sub> is hydrogen. Preferred compounds of formula I that are useful for preparing compounds of formula I  
5 wherein R<sub>2</sub> is hydrogen are compounds wherein R<sub>2</sub> is methyl, ethyl, propyl, isopropyl, acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, benzyl, or *p*-toluenesulfonyl.

The invention also provides a method for preparing a compound of formula (I) wherein R<sub>2</sub> is hydrogen comprising deprotecting a corresponding  
10 compound of formula (I) wherein R<sub>2</sub> is a suitable nitrogen protecting group.

Compounds of the invention can generally be prepared using the synthetic schemes illustrated in Figures 1-6. Starting materials can be prepared by procedures described in these schemes or by procedures that would be well known to one of ordinary skill in organic chemistry. The variables used in the  
15 Schemes are as defined below or as in the claims.

Compounds of formula I can be prepared by reactions outlined in Figure 1. Step 1 involves formation of intermediate N-nitroso compounds by treatment with isoamylnitrite or other standard acid catalyzed N-nitrosation conditions. The resulting N-nitroso compounds are directly reduced to their  
20 corresponding hydrazines (2) by treatment with lithium aluminum hydride in a suitable solvent such as tetrahydrofuran. In step 2, hydrazines (2) are condensed by acid catalysis with 1-benzoylhexahydroazepine, which is available by the process described in *J. Org. Chem.*, Vol. 33, pp 3187-95 (1968), or with benzyl 4-oxo-1-azepanecarboxylate, the synthesis of which is described in the  
25 experimental section. Fischer/Indole cyclization of the crude hydrazines provides the desired azepinoindoles (3). The Fischer/Indole cyclization can be effected with a variety of acids such as formic acid, acetic acid, trifluoroacetic acid, aqueous hydrochloric acid, aqueous sulfuric acid, or polyphosphoric acid. Step 3 is effected by either catalytic hydrogenolysis when R<sub>2</sub> is benzyl or  
30 benzyloxycarbonyl, or by base catalyzed hydrolysis in a suitable solvent such as

ethylene glycol when R<sub>2</sub> is benzoyl. Azepinoindoles 4 (wherein when R<sub>2</sub> is hydrogen) can conveniently be isolated as their hydrochloride salts.

It will be apparent to those skilled in the art that many of the requisite amines (1) are commercially available or known in the literature. The  
5 necessary 3,4-dihydro-1(2H)-quinolinyamines required for Examples 1-13 are known compounds. Indolines required for the synthesis of Examples 14-18 are either commercially available or readily prepared from known indoles following the procedure described in *Synthesis* pp. 859-60 (1977). For the  
10 benzmorpholines and benzthiomorpholines required for Examples 19-25, the synthetic route shown in Figure 2 was followed. Nitrophenols 5 were alkylated with ethyl bromoacetate derivatives to afford 6. The nitro moiety is then reduced with Pd/C in the presence of hydrogen in a suitable solvent such as ethanol. In situ cyclization gives amide 7, which is then reduced with borane to provide the required amines 8.

15 Compounds of formula I can also be prepared by the reactions outlined in Figure 3. Azapinoindoles 9 are known in the literature (*J Med Chem.*, 1968, 11, 101-106) and can participate in a Michael conjugate addition into ethyl acrylate, or a derivative thereof, in the presence of a suitable base such as cesium carbonate. Base-catalyzed hydrolysis of esters 10 gives crude acids  
20 which then undergo intramolecular Friedel-Crafts acylation in an acidic media (e.g. polyphosphoric acid or Eaton's reagent). When R<sub>2</sub> is benzoyl, azepinoindoles 12 can be obtained from base-catalyzed hydrolysis in tetrahydrofuran/methanol.

Aryl ketones 11 can be used as intermediates in the synthesis of  
25 additional compounds of formula I as shown in Figure 4. The ketone moiety is reduced with sodium borohydride to give alcohols 13. When R<sub>2</sub> is benzoyl, base-catalyzed hydrolysis of alcohols 13 gives azepinoindoles 14. Alternatively, alcohols 13 can also be used as intermediates in the synthesis of additional compounds of formula I as shown in Figure 5. Alcohols 13 can be alkylated with an alkyl halide  
30 in the presence of sodium hydride or with phenols via Mitsunobu reaction

conditions. The use of thiols or amines in the Mitsunobu reaction can give additional derivatives. Removal of  $R_2$  gives azepinoindoles 16.

Azepinoindoles 9 can also be used to make compounds of formula I as shown in Figure 6. Alkylation of 9 with 2-bromomethyl-1,3-dioxolane, or a derivative thereof, gives compounds 17. Acid-catalyzed removal of the acetal group leads to an aldehyde that can be reacted with trimethylsulfoxonium iodide and sodium hydride to give epoxides 18. Use of a Lewis acid, such as boron trifluoride diethyl etherate, would lead to alcohols 19. These alcohols could be treated as above to give azepinoindoles 20 and 21.

10 In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compounds as salts may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, 15 succinate, benzoate, ascorbate,  $\alpha$ -ketoglutarate, and  $\alpha$ -glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard 20 procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

25 Compounds of the present invention can conveniently be administered in a pharmaceutical composition containing the compound in combination with a suitable excipient. Such pharmaceutical compositions can be prepared by methods and contain excipients which are well known in the art. A generally recognized compendium of such methods and ingredients is

Remington's Pharmaceutical Sciences by E.W. Martin (Mark Publ. Co., 15th Ed., 1975). The compounds and compositions of the present invention can be administered parenterally (for example, by intravenous, intraperitoneal or intramuscular injection), topically, orally, or rectally.

5                   For oral therapeutic administration, the active compound may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course,  
10 be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

                  The tablets, troches, pills, capsules, and the like may also contain  
15 the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring  
20 may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and  
25 the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the active  
30 compound may be incorporated into sustained-release preparations and devices.

The compounds or compositions can also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid  
5 polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

Pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the  
10 active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example,  
15 water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action  
20 of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of  
25 agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile  
30 injectable solutions, the preferred methods of preparation are vacuum drying and



the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the present compounds may be applied  
5 in pure form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid  
10 carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the present compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent  
15 pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers. Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for  
20 application directly to the skin of the user.

Useful dosages of the compounds of formula I can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

25 The compound is conveniently administered in unit dosage form; for example, containing about 0.05 mg to about 500 mg, conveniently about 0.1 mg to about 250 mg, most conveniently, about 1 mg to about 150 mg of active ingredient per unit dosage form. The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate  
30 intervals, for example, as two, three, four or more sub-doses per day. The sub-

dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations.

The compositions can conveniently be administered orally, sublingually, transdermally, or parenterally at dose levels of about 0.01 to about 150 mg/kg, preferably about 0.1 to about 50 mg/kg, and more preferably about 0.1 to about 10 mg/kg of mammal body weight.

For parenteral administration the compounds are presented in aqueous solution in a concentration of from about 0.1 to about 10%, more preferably about 0.1 to about 7%. The solution may contain other ingredients, such as emulsifiers, antioxidants or buffers.

The exact regimen for administration of the compounds and compositions disclosed herein will necessarily be dependent upon the needs of the individual subject being treated, the type of treatment and, of course, the judgment of the attending practitioner.

The ability of a compound of the invention to act as a 5-HT receptor agonist or antagonist can also be determined using *in vitro* and *in vivo* assays that are known in the art. The invention provides compounds of formula I that act as either agonists or as antagonists of one or more 5-HT receptor subtypes. The compounds Exemplified herein are 5-HT ligands, which typically displace >50% of a radiolabeled test ligand from one or more 5-HT receptor subtype at a concentration of 1 M. The procedures used for testing such displacement are well known and would be readily available to one skilled in the art.

## DESCRIPTION OF PREFERRED EMBODIMENTS

### PREPARATION 1.

#### Preparation of 1-benzyl 4-ethyl 5-oxo-1,4-azepanedicarboxylate


A dry 500 ml 3-neck flask was charged with benzyl 4-oxo-1-

piperidinecarboxylate (35.08 g, 150 mmol). It was dissolved in 130 ml Et<sub>2</sub>O and cooled to -45°C. Ethyldiazoacetate (20.5 ml, 195 mmol) and boron trifluoride ethyl ether (19.4 ml, 158 mmol) were added simultaneously by syringe pump over 45 minutes. The temperature was kept below -25°C. The reaction was stirred for 30 minutes longer, and then quenched with sat. NaHCO<sub>3</sub>. The ice bath was removed. The reaction was diluted with EtOAc (250 ml) and H<sub>2</sub>O (150 ml). The layers were separated, and the organic phase was dried over MgSO<sub>4</sub>. It was concentrated under reduced pressure to an orange oil. The product was purified by flash chromatography (silica gel, 40% EtOAc/hexane), yielding product as pale yellow oil (42.1 g, 88%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.39-7.31, 5.15-5.12, 4.42-4.17, 3.96-3.83, 3.75-3.70, 3.65, 3.54-3.37, 2.08-2.03, 1.29-1.24; IR (liq.) 1743, 1702, 1476, 1455, 1443, 1425, 1371, 1318, 1295, 1238, 1213, 1178, 1098, 1068, 1028 cm<sup>-1</sup>.

#### Preparation of benzyl 4-oxo-1-azepanecarboxylate

A solution of potassium hydroxide (24.6 g, 375 mmol) in H<sub>2</sub>O (400 ml) was added to a solution of 1-benzyl 4-ethyl 5-oxo-1,4-azepanedicarboxylate (40.0 g, 125 mmol) in ethanol (400 ml). The resulting mixture was heated at reflux for 2.5 hours. Reaction was then cooled to rt., the ethanol was removed under reduced pressure, and was diluted with 200 ml brine and 300 ml ethyl acetate. The layers were separated, and the aqueous phase was extracted with ethyl acetate (2x 100 ml). The combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, and concentrated to an orange oil *in vacuo*. The product was isolated by flash chromatography (silica gel, 40%EtOAc/hexane) yielding a clear, colorless oil (22.6 g, 73%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.31-7.30, 5.12, 3.65-3.63, 2.68-2.60, 1.81-1.78; IR (liq.) 1698, 1475, 1454, 1442, 1423, 1331, 1320, 1295, 1270, 1241, 1191, 1165, 1091, 900, 699 cm<sup>-1</sup>.

EXAMPLE 1 Preparation of 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride



5

Step 1. Preparation of 3,4-dihydro-1(2H)-quinolinylamine hydrochloride.

A neat reaction between 1,2,3,4-tetrahydroquinoline (3.71 g, 27.9 mmol) and isoamyl nitrite (8.72 g, 74.4 mmol) was allowed to stir for 1 hour.

- 10 The residual isoamyl nitrite was removed under reduced pressure and the N-nitroso intermediate was taken up in tetrahydrofuran (20 ml). This solution was added dropwise to a refluxing solution of lithium aluminum hydride in tetrahydrofuran (55 ml, 1M, 55.0 mmol). One hour after the addition was complete, the reaction was cooled to 0 °C and quenched. The reaction was
- 15 filtered, concentrated under reduced pressure, and extracted into ether. The ethereal solution was washed with water, brine and dried over anhydrous potassium carbonate. The resulting brown oil (4.11 g) was trapped as the hydrochloride salt and recrystallized from methanol\ethyl acetate\hexanes to provide the title compound (mp 186-189 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.55, 7.31,
- 20 7.10, 6.98, 6.87, 3.59, 2.74, 2.05;  
IR (drift) 3053, 2956, 2940, 2868, 2841, 2835, 2724, 2667, 1603, 1586, 1580, 1567, 1545, 1499, 747 cm<sup>-1</sup>.

- Step 2. Preparation of benzyl 5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate.
- 25

A solution of 3,4-dihydro-1(2H)-quinolinylamine (2.97 g, 20.1 mmol), benzyl 4-oxo-1-azepanecarboxylate (4.96 g, 20.1 mmol), and glacial acetic acid (0.2 ml) in ethanol (200 ml) was allowed to reflux for 2.5 hours. The reaction was then cooled and evaporated *in vacuo*. The hydrazone product was purified by flash chromatography (90g SiO<sub>2</sub>, 1% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) providing benzyl 4-[3,4-dihydro-1(2H)-quinolinylimino]-1-azepanecarboxylate (7.55 g) as an oil. To a solution of benzyl 4-[3,4-dihydro-1(2H)-quinolinylimino]-1-azepanecarboxylate (6.79g, 17.99 mmol) in ethanol (200 ml) was added trifluoroacetic acid (6.22 g, 53.96 mmol). The reaction was heated and allowed to stir at reflux for 2.5 hours, at which time it was cooled to rt., evaporated, and extracted into dichloromethane. This extract was washed with water, brine, dried with anhydrous sodium sulfate, and evaporated under reduced pressure. Crystallization from ethyl acetate/hexanes provided 2.93 g of the title compound (mp 131-133 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.38, 7.29, 7.00, 6.86, 5.19, 3.98, 3.78, 2.97, 2.24; IR (drift) 2947, 1699, 1473, 1420, 1358, 1260, 1250, 1235, 1216, 1101, 997, 764, 757, 746, 703 cm<sup>-1</sup>.

Step 3. Preparation of 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride.


A mixture of benzyl 5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate (2.42 g, 6.71 mmol) and 10 % Pd/C (0.15 g) in ethanol (110 ml) was hydrogenated under 40 psi for 1.5 hours. The mixture was filtered through celite, rinsed with methanol, dichloromethane and evaporated. Methanolic hydrochloric acid is added and evaporated. The resulting product is recrystallized from methanol/ethyl acetate to give 1.40 g (80%) of the title compound (mp 261-263 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.25, 6.95, 6.84, 4.08, 3.48, 3.32, 3.28, 3.20, 2.95, 2.22; IR (drift) 2939, 2894,

2880, 2863, 2832, 2806, 2759, 2743, 2687, 2669, 2645, 2445, 1332, 1253, 743  
cm<sup>-1</sup>.

**EXAMPLE 2**

Preparation of 2-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride

5




Following the general procedure of Example 1, making non-critical  
variations but starting with 6-methyl-1,2,3,4-tetrahydroquinoline, the title  
compound was obtained (mp 268-270 °C). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 9.6, 6.99, 6.61,  
3.98, 3.30, 3.16, 3.03, 2.81, 2.32, 2.08; IR (drift) 2965, 2949, 2934, 2902, 2880,  
2850, 2823, 2791, 2769, 2695, 2659, 2438, 1458, 1251, 839 cm<sup>-1</sup>.

15

**EXAMPLE 3**

Preparation of 1-methoxy-5,6,9,10,11,12-hexahydro-  
4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline  
hydrochloride




20

Following the general procedure of Example 1, making non-critical  
variations but starting with 7-methoxy-1,2,3,4-tetrahydroquinoline, the title


compound was obtained (mp 277-279 °C). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 9.27, 6.65, 6.32, 3.96, 3.77, 3.29, 3.15, 2.78, 2.05; IR (drift) 2968, 2953, 2935, 2889, 2832, 2803, 2780, 2765, 2744, 2713, 1594, 1509, 1247, 1152, 781 cm<sup>-1</sup>.

5    **EXAMPLE 4**                      Preparation of 2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride



10                      Following the general procedure of Example 1, making non-critical variations but starting with 6-fluoro-1,2,3,4-tetrahydroquinoline, the title compound was obtained (mp 273-276 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.94, 6.64, 4.07, 3.50, 3.44, 3.32, 3.28, 3.15, 2.94, 2.22; IR (drift) 2957, 2926, 2894, 2843, 2814, 2799, 2742, 2708, 2665, 2561, 2545, 2440, 1496, 1419, 1129 cm<sup>-1</sup>.

15                      **EXAMPLE 5**                      Preparation of 6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride




20

Following the general procedure of Example 1, making non-critical variations but starting with 2-methyl-1,2,3,4-tetrahydroquinoline, the title compound is obtained (mp 214-217 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.61, 7.25, 6.93, 6.84,


4.62, 3.32, 3.11, 2.87, 2.12, 2.01, 1.20; IR (drift) 2968, 2933, 2893, 2814, 2742, 2711, 2669, 2560, 2437, 1463, 1415, 1377, 1335, 1324, 742  $\text{cm}^{-1}$ .

EXAMPLE 6 Preparation of 5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride



Following the general procedure of Example 1, making non-critical variations but starting with 3-methyl-1,2,3,4-tetrahydroquinoline, the title compound was obtained (mp 273-274 °C).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  10.10, 7.27, 7.02, 6.88, 4.07, 3.51, 3.37, 3.31, 3.00, 2.65, 2.37, 1.20; IR (drift) 2956, 2923, 2899, 2867, 2831, 2800, 2766, 2686, 2670, 2562, 2448, 1454, 1428, 1257, 740  $\text{cm}^{-1}$ .

EXAMPLE 7 Preparation of 4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride




Following the general procedure of Example 1, making non-critical variations but starting with 4-methyl-1,2,3,4-tetrahydroquinoline, the title compound was obtained (mp 260-263 °C).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  9.79, 7.22, 6.92,



6.85, 4.10, 4.00, 3.28, 3.20, 3.08, 2.15, 1.80, 1.30; IR (drift) 2959, 2939, 2891, 2873, 2860, 2811, 2800, 2739, 2709, 2662, 2647, 2553, 2542, 2437, 734  $\text{cm}^{-1}$ .

EXAMPLE 8 Preparation of (+)-6-methyl-5,6,9,10,11,12-hexahydro-  
4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline  
hydrochloride




10

Following the general procedure of Example 5 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase was 5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound was obtained (mp 196-199 °C).

EXAMPLE 9 Preparation of (-)-6-methyl-5,6,9,10,11,12-hexahydro-  
4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline  
hydrochloride

20

32




Following the general procedure of Example 5 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase was 5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound was obtained (mp 196-199 °C);  $[\alpha]_D^{25} = -27$  (c 0.88, DMSO).

10

EXAMPLE 10 Preparation of (+)-5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride

15




Following the general procedure of Example 6 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The

20

column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase was 5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound was obtained (mp 259-262 °C);  $[\alpha]_D^{25} = +20$  (c 0.28, DMSO).

5

EXAMPLE 11 Preparation of (-)-5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride




10

Following the general procedure of Example 6 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase is 5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound was obtained (mp 259-262 °C);  $[\alpha]_D^{25} = -21^\circ$  (c 0.43, DMSO).

15

20 EXAMPLE 12 Preparation of (+)-4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride

34




(+)

Following the general procedure of Example 7 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase was 5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound was obtained (mp 261-263 °C);  $[\alpha]_D^{25} = +39$  (c 0.41, chloroform).

10

**EXAMPLE 13** Preparation of (-)-4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline hydrochloride




(-)

15

Following the general procedure of Example 7 making non-critical variations, preparative chiral HPLC was performed after step two on an EM ST 140R closed loop recycling prep HPLC system (EM Separations Technology). The column used was a 5x50 cm Chiralpak AD column at 30 °C. The mobile phase was

5% isopropanol/95% heptane at a flow rate of 75 ml/min. Peak collection was monitored by UV detection at 285 nm. Following step three, the title compound is obtained (mp 261-263 °C);  $[\alpha]_D^{25} = -39$  (c 0.51, chloroform).

5    EXAMPLE 14        Preparation of 4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole hydrochloride




10                      Following the general procedure outlined in EXAMPLE 1, starting with indoline, and utilizing 1-benzoylhexahydroazepine as the ketone in step 4, 4,5,8,9,10,11-hexahydro-9H-azepino[4,5-b]pyrrolo[3,2,1-hi]indol-9-yl(phenyl)methanone was obtained (mp 130-133 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.42, 7.08, 6.94, 6.85, 4.28, 4.00, 3.76, 3.63, 3.15, 2.88; MS (ESI+) for C<sub>21</sub>H<sub>20</sub>N<sub>2</sub>O H  
15    *m/z* 240.1 (M+H)<sup>+</sup>.

Step 4        A mixture of 4,5,8,9,10,11-hexahydro-9H-azepino[4,5-b]pyrrolo[3,2,1-hi]indol-9-yl(phenyl)methanone (1.0 g, 3.2 mmol) and potassium hydroxide (0.89 g, 15.8 mmol) in ethylene glycol (10 ml) was heated under N<sub>2</sub> at  
20    170 °C for 3 h. The reaction was cooled to rt, poured into water (50 ml) and extracted with methylene chloride (4 x 50 ml). The combine organics were washed with brine, dried over anhydrous potassium carbonate, and concentrated *in vacuo*.

The residue was trapped as its HCl salt by treatment of a solution of the product in methanol with ethereal HCl. The resulting precipitate is recrystallized from  
25    methanol and ethyl acetate to give the desired product (0.4 g) (mp 249-250 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.16, 6.93, 6.82, 4.38, 3.74, 3.08, 2.94; IR (drift) 2925, 2916, 2885, 2847, 2805, 1510, 1462, 1409, 1350, 1336, 1306, 1279, 767, 758, 746 cm<sup>-1</sup>.

EXAMPLE 15 Preparation of 2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole hydrochloride




5

Following the general procedure outlined in EXAMPLE 14, and making non-critical variations but starting with 5-fluoro indoline, the title compound was obtained (mp 250-252 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.68, 6.43, 4.15, 3.47, 2.88, 2.76; IR (drift) 2930, 2911, 1661, 1507, 1412, 1354, 1261, 1171, 1112, 938, 857, 848, 839, 708, 688 cm<sup>-1</sup>.

10

EXAMPLE 16 Preparation of 2-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole hydrochloride




15

Following the general procedure outlined in EXAMPLE 14, and making non-critical variations but starting with 5-methoxy indoline, the title compound was obtained (mp 269-271 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.70, 6.57, 4.44, 3.80, 3.72, 3.45, 3.23, 3.16; IR (drift) 2974, 2948, 2908, 2891, 2841, 2817, 2805, 2769, 2718, 1509, 1421, 1255, 1244, 1233, 1142 cm<sup>-1</sup>.

20


EXAMPLE 17 Preparation of 5-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole hydrochloride

37




Following the general procedure outlined in EXAMPLE 1, and making non-critical variations but starting with 2-methyl indoline, the title compound was obtained (mp 243-246 °C). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.13, 6.91, 7.80, 4.99-4.87, 3.93, 3.51-3.23, 3.17, 1.47; IR (drift) 2973, 2960, 2941, 2927, 2901, 2883, 2848, 2826, 2794, 2737, 2650, 2549, 2438, 1292, 751 cm<sup>-1</sup>.

EXAMPLE 18 Preparation of 4,5-dimethyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole hydrochloride




Following the general procedure outlined in EXAMPLE 1, and making non-critical variations but starting with 2, 3-dimethyl indole, the title compound was obtained (mp 195-197 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.16, 6.92, 6.81, 4.40, 3.64, 3.53-3.35, 3.16, 1.54, 1.43; IR (drift) 2960, 2938, 2925, 2866, 2847, 2821, 2727, 2657, 2635, 2533, 2427, 1465, 1372, 1287, 746 cm<sup>-1</sup>.

EXAMPLE 19 Preparation of 2,3,4,5,8b,9,10,11,12,12a-decahydro-1H-azepino[4',5':4,5]pyrrolo[3,2,1-jk]carbazole hydrochloride



Following the general procedure outlined in EXAMPLE 1, and making non-critical variations but starting with 1,2,3,4 tetramethyl carbazole, the title compound was obtained (mp 210-212 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.16, 6.94, 6.82, 4.75, 4.14, 3.42, 3.25, 3.16, 2.12-1.96, 1.54, 1.41, 1.26, 1.08; IR (drift) 3046, 2936, 2929, 2846, 2810, 2744, 2633, 2541, 2519, 2429, 1460, 1328, 1290, 754, 744 cm<sup>-1</sup>;

EXAMPLE 20 Preparation of 6-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride



Step 5. Preparation of 6-methyl-2H-1,4-benzoxazin-3(4H)-one

To a solution of 4-methyl-2-nitrophenol (9.95 g, 65.0 mmol) in acetone (170 ml) was added, in one portion, potassium carbonate (10.8 g, 78.0 mmol) and ethyl bromoacetate (7.9 ml, 71.5 mmol). The resulting mixture was heated at reflux for 3.5 hours. The acetone was removed *in vacuo*, and the resulting material was partitioned between ethyl acetate and water. The layers were separated, and the aqueous phase was extracted twice with ethyl acetate. The ethyl acetate layer was washed with saturated sodium bicarbonate, brine, dried over magnesium sulfate, filtered, and evaporated to afford an orange oil. The intermediate nitro compound was taken up in ethanol (150 ml) and hydrogenated with Pd/C (0.33 g) at 51 psi for 90 minutes. The mixture was filtered through celite, and the filtrate evaporated. The product was recrystallized from methanol to yield




7.29 g of the title compound as an off-white solid (mp 204-205 °C). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 6.81-6.78, 6.70-6.67, 4.48, 2.18; MS [MH<sup>+</sup>] 162.1.

Step 6. Preparation of 6-methyl-3,4-dihydro-2H-1,4-benzoxazine

To a solution of 6-methyl-2H-1,4-benzoxazin-3(4H)-one (7.02 g, 43.0 mmol) in tetrahydrofuran (130 ml) was added borane dimethylsulfide complex (43 ml, 10M). This solution was stirred at rt for 2.5 hours, then quenched with 1 M hydrochloric acid. The solvent was evaporated, and the residue was partitioned between saturated sodium bicarbonate and dichloromethane. The layers were separated, and the aqueous phase was extracted twice with dichloromethane. The dichloromethane layer was washed with brine, dried over magnesium sulfate, and evaporated to yield 5.1 g of the title compound as a pale yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.70-6.68, 6.49-6.46, 6.42, 4.24-4.21, 3.41-3.38, 2.22; MS [MH<sup>+</sup>] 150.2.

Following the remainder of the general procedure of Example 1 (steps 1-3) and making non-critical variations but starting with 6-methyl-3,4-dihydro-2H-1,4-benzoxazine, the title compound was obtained (mp 271-273 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.59, 6.40, 4.44, 4.17, 3.52, 3.49, 3.27, 3.24, 2.57; IR (drift) 2979, 2957, 2934, 2872, 2855, 2828, 2792, 2760, 2746, 2683, 2652, 1511, 1265, 1243, 1223 cm<sup>-1</sup>.

EXAMPLE 21 Preparation of 1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride




Following the general procedure of Example 20 but starting with 2

H-1,4-Benzoxazine-3(4 H)-one, utilizing 1-benzoylhexahydroazepine as the ketone in the hydrazone formation of step two, and thus the deprotection sequence described in step 4 of example 14, the title compound was obtained (mp 217-219 °C). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 9.6, 7.01, 6.85, 6.50, 4.46, 4.18, 3.28-3.08, 2.50; IR (drift) 2949, 2925, 2878, 2843, 2812, 2758, 2689, 2673, 1499, 1328, 1247, 1036, 872, 771, 729 cm<sup>-1</sup>.

EXAMPLE 22 Preparation of 6-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride


10



Following the general procedure of Example 20 and making non-critical variations but starting with 2-amino-4-chlorophenol, the title compound was obtained (mp decompose >275°C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.84, 6.50, 4.49-4.48, 4.22-4.20, 3.65-3.63, 3.53-3.48, 3.30-3.27; IR (free amine) (drift) 2932, 2897, 2882, 2823, 1495, 1466, 1379, 1355, 1323, 1277, 1269, 1236, 1217, 1199, 1014 cm<sup>-1</sup>.


EXAMPLE 23 Preparation of 5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride

20




Following the general procedure of Example 20 and making non-critical variations but starting with 5-fluoro-2-nitrophenol, the title compound was obtained (mp decompose  $>250^{\circ}\text{C}$ ).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.76-6.72, 6.38-6.33, 4.52-4.49, 4.20-4.17, 3.51-3.43, 3.28-3.25, 3.18-3.14; IR (drift) 2950, 2855, 2805, 2767, 2757, 2710, 2675, 2651, 2565, 2449, 1645, 1591, 1500, 1333, 1109  $\text{cm}^{-1}$ .

EXAMPLE 24 Preparation of 5-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride



Following the general procedure of Example 20 but starting from 5-methyl-2-nitrophenol and using glacial acetic acid for the cyclization in step 3, the title compound was obtained (mp  $274-275^{\circ}\text{C}$ ).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.81, 6.37, 4.44, 4.12, 3.48-3.41, 3.22, 3.15, 2.36; IR (drift) 2983, 2965, 2940, 2880, 2856, 2822, 2809, 2758, 2731, 2667, 1590, 1503, 1331, 1031, 847  $\text{cm}^{-1}$ .


EXAMPLE 25 Preparation of 6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride



Following the general procedure of Example 20 but starting from 4-


fluoro-2-nitrophenol and using *p*-toluene sulfonic acid for the cyclization in step 3, the title compound was obtained (mp 257-259°C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.55-6.49, 6.43-6.40, 4.44, 4.17, 3.49-3.44, 3.38-3.33, 3.26-3.23; IR (drift) 2949, 2876, 2845, 2815, 2768, 2685, 2676, 2626, 1511, 1362, 1277, 1224, 1023, 881, 791 cm<sup>-1</sup>.

**EXAMPLE 26** Preparation of 2-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride




Following the general procedure of Example 20 and making non-critical variations but starting with 2-nitrophenol and ethyl-2-bromopropionate, the title compound was obtained (mp 255-257 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.01, 6.88, 6.54, 4.40-4.35, 4.30, 3.68-3.63, 3.49-3.42, 3.25-3.17, 1.51; IR (drift) 2967, 2931, 2809, 2790, 2736, 2711, 2655, 2647, 2557, 2439, 1502, 1377, 1322, 1243, 794 cm<sup>-1</sup>.

**EXAMPLE 27** Preparation of 1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]thiazino[2,3,4-hi]indole hydrochloride




Following the general procedure of Example 21 but starting from 2H-1,4-benzothiazin-3(4H)-one and using 10% sulfuric acid for the cyclization in step 3, the title compound was obtained (mp 263 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.24, 6.94, 6.87, 3.36-3.24, 3.12-3.10; IR (drift) 2937, 2929, 2862, 2814, 2718, 2663, 2626, 2594, 2544, 2428, 1463, 1412, 1335, 782, 739 cm<sup>-1</sup>.

EXAMPLE 28 Preparation of 2,2-dimethyl-1,2,8,9,10,11-hexahydro-7H-azepino [4,5-b] [1,4]oxazino[2,3,4-hi]indole hydrochloride



Following the general procedure of Example 20 but starting with 2-nitrophenol and ethyl-2-bromoisobutyrate, and using p-toluene sulfonic acid for the cyclization in step 3, the title compound was obtained (mp decompose >260°C). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.03, 6.92, 6.52, 3.96, 3.53-3.46, 3.26-3.20, 1.41; IR (drift) 2972, 2948, 2858, 2824, 2766, 2751, 1583, 1498, 1385, 1332, 1245, 1202 cm<sup>-1</sup>.


EXAMPLE 29 Preparation of 4-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole maleic acid



Following the general procedure of Example 20 but starting with 2-fluoro-6-nitrophenol, using p-toluene sulfonic acid for the cyclization in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 176-


177°C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.95, 6.78, 6.24, 4.51, 4.19, 3.42-3.50, 3.23, 3.16; IR (drift) 3009, 2925, 2899, 2879, 2835, 2786, 2714, 1705, 1626, 1550, 1531, 1526, 1486, 1358  $\text{cm}^{-1}$ .

5    **EXAMPLE 30**            Preparation of 4-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole maleic acid



Following the general procedure of Example 20 but starting with 6-chloro-2-nitrophenol, using p-toluene sulfonic acid for the cyclization in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 183-184 °C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.99, 6.91, 6.24, 4.56, 4.20, 3.43-3.51, 3.23, 3.16; IR (drift) 1645, 1639, 1627, 1608, 1568, 1536, 1517, 1497, 1476, 1385, 1377, 1372, 1355, 1196  $\text{cm}^{-1}$ .


15    **EXAMPLE 31**            Preparation of 5-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole maleic acid



Following the general procedure of Example 20 but starting with 5-chloro-2-nitrophenol, using TFA in EtOH for the cyclization in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 177-179 °C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  7.05, 6.55, 6.24, 4.50, 4.18, 3.42-3.50, 3.23-3.25, 3.14-3.16;


IR (drift) 2892, 2867, 2779, 2745, 1627, 1569, 1536, 1491, 1466, 1451, 1369, 1329, 1024  $\text{cm}^{-1}$ .

EXAMPLE 32 Preparation of 6-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride



Following the general procedure of Example 20 but starting with 2-nitro-4-(trifluoro methyl)-phenol, using p-toluene sulfonic acid for the cyclization in step 3, the title compound was obtained (mp 248-250 °C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  7.30, 6.61, 4.55, 4.26, 3.46-3.53; IR (drift) 2815, 2797, 2741, 2670, 2560, 1583, 1333, 1244, 1198, 1158, 1107, 1094, 1060, 1021  $\text{cm}^{-1}$ .


EXAMPLE 33 Preparation of 5,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole maleic acid



Following the general procedure of Example 20 but starting with 4,5-difluoro-2-nitrophenol, using 10% sulfuric acid in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 184-185 °C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.45, 6.23, 4.46, 4.17, 3.45-3.50, 3.34-3.37, 3.220-3.24; IR (drift) 3070,


2879, 2780, 2728, 1601, 1575, 1518, 1482, 1457, 1373, 1353, 1171, 1042, 1000  $\text{cm}^{-1}$ .

EXAMPLE 34 Preparation of 5-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole maleic acid



Following the general procedure of Example 20 but starting with 5-chloro-4-fluoro-2-nitrophenol, using 10% sulfuric acid in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 194-195 °C).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  6.52, 6.24, 4.46, 4.18, 3.51-3.45, 3.29-3.37, 3.21-3.25; IR (drift) 1615, 1583, 1552, 1501, 1480, 1456, 1368, 1355, 1280, 1222, 1169  $\text{cm}^{-1}$ .

EXAMPLE 35 Preparation of 5-fluoro-6-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole maleic acid




Following the general procedure of Example 20 but starting with 4-chloro-5-fluoro-2-nitrophenol, using 10% sulfuric acid in step 3, and preparing the Maleic acid salt, the title compound was obtained (mp 173-175 °C).  $^1\text{H}$  NMR




(CD<sub>3</sub>OD)  $\delta$  6.50, 3.25, 4.50, 4.19, 3.56-3.59, 3.44-3.50, 3.22-3.25; IR (drift) 3062, 3028, 2971, 2897, 1626, 1608, 1585, 1498, 1463, 1372, 1365, 1352, 1153, 1036 cm<sup>-1</sup>.

- 5    **EXAMPLE 36**      Preparation of 5,6-dichloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole maleic acid



- Following the general procedure of Example 20 but starting with 4,5-dichloro-2-nitrophenol, using 10% sulfuric acid in step 3, and preparing the  
 10    Maleic acid salt, the title compound was obtained (mp 180-181 °C). <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  6.68 (s, 1 H), 6.23 (s, 2 H), 4.49 (t, *J* = 4.4 Hz, 2 H), 4.20 (t, *J* = 4.8 Hz, 2 H), 3.59-3.62 (t, *J* = 2 Hz, H), 3.45-3.51 (m, 4 H), 3.22-3.26 (m, 2 H); IR (drift) 2887, 2840, 1625, 1608, 1578, 1551, 1486, 1458, 1372, 1351, 1273, 1023 cm<sup>-1</sup>.


- 15    **EXAMPLE 37**      Preparation of 4,6-dichloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole hydrochloride



- Following the general procedure of Example 20 but starting with 3,5-dichloro-2-nitrophenol, and using 10% sulfuric acid in step 3, the title  
 20    compound was obtained (decompose >255°C). <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  6.89, 4.55, 4.21, 3.56-3.59, 3.44-3.51, 3.23-3.27; IR (drift) 2955, 2895, 2743, 2652, 2635,

2556, 2428, 1494, 1468, 1420, 1345, 1294, 1229  $\text{cm}^{-1}$ .

EXAMPLE 38 Preparation of 1-chloro-2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole maleic acid




5

Following the general procedure outlined in EXAMPLE 14, and making non-critical variations but starting with 6-chloro-5-fluoro indoline, the title compound was obtained (amorphous solid).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ) 6.70, 6.23, 4.52, 3.69, 3.34-3.22, 3.08; IR (drift) 2420 (b), 1628, 1561, 1506 (s), 1449, 1421, 1378 (s), 1350, 1337, 1323, 1300, 1291, 1278, 1254, 1147  $\text{cm}^{-1}$ ; MS (ESI+) for  $\text{C}_{14}\text{H}_{14}\text{ClFN}_2$   $m/z$  265.2 ( $\text{M}+\text{H}$ ) $^+$ .

EXAMPLE 39 Preparation of 2-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole maleic acid


15



Following the general procedure outlined in EXAMPLE 14, and making non-critical variations but starting with 5-chloro-indoline, the title compound was obtained (amorphous solid).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ) 7.16, 6.91, 6.81, 4.45, 3.74, 3.40, 3.24, 3.15; MS (ESI+) for  $\text{C}_{14}\text{H}_{13}\text{ClN}_2$   $m/z$  247.2 ( $\text{M}+\text{H}$ ) $^+$ .


EXAMPLE 40 Preparation of 1-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole maleic acid

49



Following the general procedure outlined in EXAMPLE 14, and making non-critical variations but starting with 6-chloro-indoline, the title compound was obtained (amorphous solid). <sup>1</sup>H NMR (CD<sub>3</sub>OD) 6.91, 6.81, 6.77, 4.90, 3.69, 3.34, 3.21; MS (ESI+) for C<sub>14</sub>H<sub>15</sub>ClN<sub>2</sub> *m/z* 247.2 (M+H)<sup>+</sup>.

EXAMPLE 41 Preparation of 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one oxalate.



Step 7. Preparation of ethyl 3-(3-benzoyl-2,3,4,5-tetrahydroazepino[4,5-b]indol-6(1H)-yl)propanoate.

15

A mixture of 3-benzoyl-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole (5.00 g, 17.2 mmol), cesium carbonate (5.61 g, 17.2 mmol), and ethyl acrylate (1.90 mL, 17.5 mmol) in acetonitrile (250 mL) was heated under N<sub>2</sub> at 50 °C for 5 h (reaction can also be run in DMF at room temperature). Cooled to room temperature, diluted with H<sub>2</sub>O, and extracted with EtOAc. The combined organic extracts were washed with saturated aqueous NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, decanted, and concentrated under reduced pressure. The crude product was purified by silica gel chromatography with EtOAc/hexanes (2:1) to give 4.82 g (72%) of the title compound as a yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.57- 7.38, 7.32, 7.23-7.07, 4.49-4.33, 4.20-4.02, 3.77-3.66, 3.28-3.13, 2.94, 2.72, 1.24; IR (liq.) 2981, 1731, 1631, 1467, 1445, 1422, 1381, 1368, 1349, 1319, 1295, 1272,

1242, 1187, 742, 707  $\text{cm}^{-1}$ ; MS (ESI) 391.0 ( $\text{M}^+ + \text{H}$ ).

Step 8. Preparation of 10-benzoyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one.

5

A solution of potassium hydroxide (0.80 g, 14.3 mmol) in  $\text{H}_2\text{O}$  (20 mL) was added to a solution of ethyl 3-(3-benzoyl-2,3,4,5-tetrahydroazepino[4,5-b]indol-6(1H)-yl)propanoate (4.25 g, 10.9 mmol) in THF (30 mL). The reaction mixture was heated at 60  $^\circ\text{C}$  for 1 h, then cooled to room temperature, acidified with 10% aqueous HCl, and extracted with EtOAc. The combined organic extracts were washed with saturated aqueous NaCl, dried over  $\text{Na}_2\text{SO}_4$ , decanted, and concentrated under reduced pressure to give 3.73 g of crude acid. The crude acid (2.00 g, 5.52 mmol) was then added to neat PPA (18.6 g) stirring at 100  $^\circ\text{C}$  under  $\text{N}_2$ . After 1.5 h, the reaction was cooled to room temperature, quenched with ice and 10% aqueous NaOH, and then extracted with EtOAc. The combined organic extracts were washed with saturated aqueous NaCl, dried over  $\text{Na}_2\text{SO}_4$ , decanted, and concentrated under reduced pressure. The crude product was purified by silica gel chromatography with EtOAc/hexanes (gradient 1:1 to 2:1) to give 0.81 g (43%) of the title compound as a yellow solid (mp 177.5-181  $^\circ\text{C}$ ).  $^1\text{H}$  NMR ( $d$ -DMSO)  $\delta$  7.75, 7.46, 7.09, 4.45-4.24, 3.93, 3.61, 3.34, 3.17, 3.11-2.80; IR (drift) 1676, 1628, 1587, 1493, 1483, 1466, 1428, 1357, 1324, 1295, 1276, 1266, 1191, 754, 706  $\text{cm}^{-1}$ .

20


Step 9. Preparation of 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one oxalate.

25

A solution of 10-benzoyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one (0.20 g, 0.58 mmol) in THF/MeOH/50% aqueous NaOH (3:2:1, 9 mL) was heated under  $\text{N}_2$  at reflux for 4 d.

Cooled to room temperature, diluted with saturated aqueous NaCl, and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic extracts were washed with saturated aqueous NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, decanted, and concentrated under reduced pressure. The crude product was purified by silica gel chromatography with  
5 CH<sub>2</sub>Cl<sub>2</sub>/MeOH/Et<sub>3</sub>NH (gradient 95:5:0 to 95:4:1) to give 0.10 g (74%) of the free base of title compound as a dark yellow foam. The oxalate salt was prepared by treating a solution of the free base in methanol with a solution of oxalic acid in ether to give a precipitate that crystallized from methanol as small yellow needles of the title compound (mp 199-202 °C [dec.]). <sup>1</sup>H NMR (*d*-  
10 DMSO) δ 7.78, 7.46, 7.15, 4.39, 4.19, 3.39, 3.34, 3.24, 3.14, 3.02; IR (drift) 2965, 2890, 2841, 2799, 2732, 2516, 2484, 1720, 1673, 1626, 1607, 1591, 1492, 1460, 1206 cm<sup>-1</sup>.


15 **EXAMPLE 42** Preparation of 2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one maleate.



20 Following the general procedure of Example 41, making non-critical variations but starting with 3-benzoyl-9-fluoro-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole and using maleic acid for the salt formation, the title compound was obtained (mp 242-244 °C [dec.]). <sup>1</sup>H NMR (*d*-DMSO) δ 9.18, 7.67, 7.21, 4.40, 3.33, 3.25, 3.12, 3.04; IR (drift) 3358, 2970, 2959, 2830, 2792, 2785, 2752, 1681, 1593, 1490, 1464, 1383, 1330, 1216, 907 cm<sup>-1</sup>.


25 **EXAMPLE 43** Preparation of 2-chloro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one oxalate.

52




Following the general procedure of Example 41, making non-critical variations but starting with 3-benzoyl-9-chloro-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole, the title compound was obtained (mp 236.5-238 °C [dec.]). <sup>1</sup>H NMR (*d*-DMSO) δ 7.89, 7.38, 4.41, 3.69, 3.38, 3.33, 3.23, 3.13, 3.04; IR (mull) 3440, 2644, 2514, 1915, 1686, 1491, 1420, 1327, 1318, 1251, 1138, 1094, 1030, 914, 720 cm<sup>-1</sup>.

EXAMPLE 44 Preparation of 6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one oxalate.




Following the general procedure of Example 41, making non-critical variations but using ethyl crotonate in step 7, the title compound was obtained (mp 215-215.5 °C). <sup>1</sup>H NMR (*d*-DMSO) δ 7.79, 7.47, 4.97, 3.50-3.20, 3.15, 2.73, 1.16; IR (drift) 3020, 2984, 2969, 1727, 1684, 1609, 1589, 1471, 1327, 1217, 1194, 1178, 1100, 752, 706 cm<sup>-1</sup>.

EXAMPLE 45 Preparation of 2-fluoro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one oxalate.




Following the general procedure of Example 41, making non-critical variations but starting with 3-benzoyl-9-fluoro-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole and using ethyl crotonate in step 7, the title compound was obtained (mp 209.5-211.5 °C). <sup>1</sup>H NMR (*d*-DMSO) δ 7.69, 7.22, 4.99, 3.44-3.22, 3.12, 2.78, 1.16; IR (drift) 2969, 1743, 1727, 1716, 1688, 1646, 1616, 1480, 1418, 1377, 1214, 1145, 1102, 858, 602 cm<sup>-1</sup>.

EXAMPLE 46 Preparation of 2,3-dichloro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one mesylate.




Following the general procedure of Example 41, making non-critical variations but starting with 3-benzoyl-8,9-dichloro-1,2,3,4,5,6-hexahydroazepino[4,5-b]indole, using ethyl crotonate in step 7, and using methanesulfonic acid for the salt formation, the title compound was obtained (mp 225-230 °C [dec.]). <sup>1</sup>H NMR (*d*-DMSO) δ 8.94, 8.12, 4.97, 3.48-3.20, 3.13, 2.77, 2.30, 1.20; IR (drift) 3029, 2979, 2849, 2792, 1683, 1481, 1410, 1312, 1214, 1196, 1184, 1161, 1145, 1040, 775 cm<sup>-1</sup>.

EXAMPLE 47 Preparation of 6-propyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one hydrochloride.



Following the general procedure of Example 41, making non-critical variations but using ethyl *trans*-hexenoate in step 7 and using hydrochloric acid for the salt formation, the title compound was obtained (mp 193-194.5 °C). <sup>1</sup>H NMR (*d*-DMSO) δ 9.38, 7.79, 7.45, 7.14, 4.87, 3.54, 3.45-  
5 3.19, 3.16, 2.83, 1.47, 1.13, 0.78; IR (drift) 2957, 2926, 2870, 2853, 2741, 1683, 1588, 1471, 1416, 1354, 1325, 1279, 1191, 795, 751 cm<sup>-1</sup>.


EXAMPLE 48 Preparation of 6-(trifluoromethyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-  
10 ij]quinolin-4-one oxalate.



Following the general procedure of Example 41, making non-critical variations but using ethyl 4,4,4-trifluorocrotonate in step 7 and using hydrochloric acid for the salt formation, the title compound was obtained (mp 201.5-202.5 °C). <sup>1</sup>H NMR (*d*-DMSO) δ 7.86, 7.54, 7.24, 5.94, 3.73, 3.67, 3.43-  
15 3.19, 3.16, 3.00; IR (drift) 1726, 1694, 1645, 1639, 1592, 1469, 1416, 1336, 1272, 1253, 1204, 1195, 1177, 1127, 945 cm<sup>-1</sup>.

20

EXAMPLE 49 Preparation of 5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol



25

Step 10 Preparation of 10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-




## 4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol.

A solution of 10-benzoyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino [4',5':4,5] pyrrolo[3,2,1-ij]quinolin-4- one (1.46 g, 4.24 mmol) in  
5 ethanol (40mL) was cooled to 0 °C under N<sub>2</sub>. Added sodium borohydride (0.32 g, 8.46 mmol) and allowed to warm slowly to room temperature. Concentrated under reduced pressure to approximately 10 mL, diluted with H<sub>2</sub>O, and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic extracts were washed with saturated aqueous NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, decanted, and concentrated under reduced  
10 pressure. The crude product was purified by silica gel chromatography with EtOAc/heptanes (2:1) to give 1.24 g (84%) of the title compound as a yellow foam. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.42, 7.33, 7.12-7.02, 5.10, 4.13, 4.02, 3.68, 3.21, 2.92, 2.38, 2.23, 1.80; IR (drift) 1627, 1614, 1475, 1466, 1457, 1448, 1429, 1372, 1360, 1329, 1292, 1268, 785, 747, 706 cm<sup>-1</sup>; MS (ESI) 369.1 (M<sup>+</sup>+Na).  
15

The 10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5] pyrrolo[3,2,1-ij]quinolin-4-ol was then hydrolyzed following the procedure outlined in step 9 to give 5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol as a tan foam. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  
20 δ 7.43, 7.09, 5.11, 4.11, 3.16, 3.02, 2.38, 2.26; IR (drift) 3300, 3047, 2923, 2878, 2828, 2751, 1479, 1454, 1430, 1415, 1371, 1330, 1199, 1074, 746 cm<sup>-1</sup>; HRMS (EI) calcd for C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O 242.1419, found 242.1422.

EXAMPLE 50      Preparation of 4-methoxy-5,6,8,9,10,11,12,12a-octahydro-  
25 4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline




Step 11. Preparation of 10-benzoyl-4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline.

NaH (60% dispersion in mineral oil, 0.50g, 1.2 mmol) was added to a solution of 10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo [3,2,1-ij]quinolin-4-ol (0.20 g, 0.58 mmol) in DMF (5.0 mL) at 0 °C under N<sub>2</sub>. After 30 min, iodomethane (0.040 mL, 0.64 mmol) was added. The reaction was quenched with saturated aqueous NH<sub>4</sub>Cl after 1 h and then allowed to warm to room temperature prior to extracting with EtOAc. The combined organic extracts were washed with saturated aqueous NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, decanted, and concentrated under reduced pressure. The crude product was purified by silica gel chromatography with EtOAc/heptanes (2:1) to give 0.16 g (79%) of the title compound as a beige foam. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.46-7.30, 6.94, 4.51, 4.22-4.03, 4.01-3.79, 3.65-3.41, 3.29, 3.12, 3.04, 2.90, 2.82, 2.34, 2.08; IR (drift) 1631, 1493, 1477, 1459, 1422, 1370, 1359, 1324, 1292, 1267, 1098, 1085, 786, 748, 706 cm<sup>-1</sup>; HRMS (FAB) calcd for C<sub>23</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>+H 361.1916, found 361.1913.

Following the general procedure of step 4, making non-critical variations but starting with 10-benzoyl-4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline, the title compound was obtained as a brown oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.45, 7.07, 4.56, 4.26, 4.09, 3.42, 3.27-3.13, 3.11-2.94, 2.49, 2.18; IR (liq.) 2926, 2905, 2882, 2820, 1492, 1479, 1453, 1415, 1370, 1332, 1200, 1098, 1083, 1066, 748 cm<sup>-1</sup>, HRMS (FAB) calcd for C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O+H 257.1654, found 257.1665.

EXAMPLE 51 Preparation of 4-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline



Step 12. Preparation of 10-benzoyl-4-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline

1,1'-Azobis(N,N-dimethylformamide) (0.34 g, 2.0 mmol) was added to a solution of 10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo [3,2,1-ij]quinolin-4-ol (0.45 g, 1.3 mmol), phenol (0.18 g, 1.9 mmol), and tributylphosphine (0.49 mL, 2.0 mmol) in dry benzene (4.0 mL) at room temperature under N<sub>2</sub>. The reaction mixture immediately congealed but resumed stirring upon heating to 60 °C. After 5 h, the reaction was cooled to room temperature and filtered to remove white precipitate. The filtrate was concentrated under reduced pressure and purified by silica gel chromatography with heptanes/EtOAc (gradient 3:1 to 3:2) to give 0.26 g (47%) of the title compound as a beige foam. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.52-7.38, 7.32, 7.11-6.95, 5.68, 4.21-3.90, 3.69, 3.21, 2.91, 2.63, 2.34; IR (drift) 1630, 1596, 1492, 1458, 1421, 1359, 1327, 1291, 1267, 1226, 1192, 786, 750, 706, 694 cm<sup>-1</sup>; MS (ESI) 423.3 (M<sup>+</sup>+H).

Following the general procedure of step 4, making non-critical variations but starting with 10-benzoyl-4-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino [4',5':4,5]pyrrolo[3,2,1-ij]quinoline, the title compound was obtained. MS (ESI) 319.2 (M<sup>+</sup>+H).

EXAMPLE 52 Using synthetic procedures similar to those described herein, the following compounds of formula (I) wherein R<sub>2</sub> is hydrogen can also be prepared:

- 4-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (100);
- 5-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (101);
- 5 6-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (102);
- 5,6-dibromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (103);
- 4,6-dibromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (104);
- 10 4-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (105);
- 5-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (106);
- 15 6-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (107);
- 4-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (108);
- 5-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (109);
- 20 4-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (110);
- 5-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (111);
- 25 6-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole (112);

- 4-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (113);
- 5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (114);
- 5 6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (115);
- 4-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (116);
- 5-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (117);
- 10 6-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (118);
- 4,5-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (119);
- 15 5,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (120);
- 4,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (121);
- 4-chloro-5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (122);
- 20 4-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (123);
- 5-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (124);
- 25 6-chloro-5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (125);

- 4-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (126);
- 5-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (127);
- 5 6-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (128);
- 4-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (129);
- 5-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (130);
- 10 6-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole; (131)
- 4-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (132);
- 15 5-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (133);
- 6-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (134);
- 4-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (135);
- 20 5-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (136);
- 6-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole (137);
- 25 1-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole (138);

- 3-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(139);
- 1-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(140);
- 5 2-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(141);
- 3-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(142);
- 3-chloro-1-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
10 (143);
- 2-chloro-1-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
hi]indole (144);
- 1-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(145);
- 15 3-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(146);
- 4-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole  
(147);
- 1-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
20 hi]indole (148);
- 2-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
hi]indole (149);
- 3-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
hi]indole (150);
- 25 4-benzyloxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (151);

- 4-(3-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (152);
- 4-(2-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (153);
- 5 4-(4-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (154);
- 4-(3-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (155);
- 4-(2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (156);
- 10 4-(4-bromo-2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (157);
- 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (158);
- 15 N-phenyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (159);
- N-(4-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (160);
- N-(3-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (161);
- 20 N-(2-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (162);
- N-(4-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (163);
- 25 N-(3-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (164);



N-(2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (165);

N-(4-bromo-2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine (166);

5 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-4-thione (167);

4-(phenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (168);

4-(4-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (169);

4-(3-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (170);

4-(2-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (171);

15 4-(4-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (172);

4-(3-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (173);

4-(2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (174);

4-(4-bromo-2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (175);

5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-ol (176);

25 5-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (177);

- 5-benzyloxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (178);
- 5-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (179);
- 5 5-(4-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (180);
- 5-(3-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (181);
- 5-(2-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (182);
- 10 5-(4-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (183);
- 5-(3-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (184);
- 15 5-(2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (185);
- 5-(4-bromo-2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (186);
- 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (187);
- 20 N-phenyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (188);
- N-(4-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (189);
- 25 N-(3-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (190);

- N-(2-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (191);
- N-(4-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (192);
- 5 N-(3-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (193);
- N-(2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (194);
- 10 N-(4-bromo-2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine (195);
- 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-5-thione (196);
- 5-(phenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (197);
- 15 5-(4-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (198);
- 5-(3-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (199);
- 5-(2-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (200);
- 20 5-(4-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (201);
- 5-(3-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (202);
- 25 5-(2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (203);

- 5-(4-bromo-2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (204);
- 4-(4-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (205);
- 5 1-[2-(4-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (206);
- 1-[2-(3-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (207);
- 1-[2-(2-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (208);
- 10 1-[2-(4-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (209);
- 1-[2-(3-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (210);
- 15 1-[2-(2-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (211);
- 1-[2-(4-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (212);
- 1-[2-(3-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (213);
- 20 1-[2-(2-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (214);
- 1-[2-(4-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (215);
- 25 1-[2-(3-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (216);

- 1-[2-(2-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (217);
- 1-[2-(4-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (218);
- 5 1-[2-(3-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (219);
- 1-[2-(2-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (220);
- 1-[2-(1-naphthyloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (221);
- 10 1-[2-(2-naphthyloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (222);
- 1-[2-([1,1'-biphenyl]-4-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (223);
- 15 1-[2-([1,1'-biphenyl]-3-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (224);
- 1-[2-([1,1'-biphenyl]-2-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (225);
- 1-[2-[4-(trifluoromethoxy)phenoxy]ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (226);
- 20 1-[2-[3-(trifluoromethoxy)phenoxy]ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (227);
- 1-[2-[2-(trifluoromethoxy)phenoxy]ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (228);
- 25 1-[2-[4-(trifluoromethyl)phenoxy]ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (229);

1-{2-[3-(trifluoromethyl)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (230); and

1-{2-[2-(trifluoromethyl)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline (231).

5

EXAMPLE 53      Using synthetic procedures similar to those described herein, the following compounds of formula (I) wherein R<sub>2</sub> is a protecting group can also be prepared:

benzyl 5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

benzyl 2-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
10-benzoyl-1-methoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-[3,2,1-ij]quinoline;

15 benzyl 2-fluoro-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

20 benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

(+)-benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

25 (-)-benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

(+)-benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

- (-)-benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(+)-benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;
- 5 (-)-benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
benzyl 2-methyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 2,2-dimethyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-
- 10 b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 4-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 4-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 15 benzyl 5-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 5-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 5-methyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-
- 20 hi]indole-9-carboxylate;  
benzyl 6-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 6-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 25 benzyl 6-methyl-1,2,6b,7,8,10,11,11a-octahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 6-(trifluoromethyl)-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 5,6-difluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-
- 30 b][1,4]oxazino[2,3,4-hi]indole-9-carboxylatebenzyl;

- benzyl 5-chloro-6-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 5-fluoro-6-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 5    benzyl 5,6-dichloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
benzyl 4,6-dichloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
7-benzoyl 4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 10    7-benzoyl 2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;  
7-benzoyl 2-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;  
benzyl-5-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]-7-
- 15    carboxylate;  
benzyl-4,5-dimethyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole-7-carboxylate;  
benzyl-2,3,4,5,8b,9,10,11,12,12a-decahydro-1H-azepino[4',5':4,5]pyrrolo[3,2,1-jk]carbazole-7-carboxylate;
- 20    benzyl-1-chloro-2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole-7-carboxylate;  
benzyl-2-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole-7-carboxylate;  
benzyl-1-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-
- 25    hi]indole-7-carboxylate;  
10-benzoyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 30    10-benzoyl-2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-2-chloro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-




- [3,2,1-ij]quinolin-4-one;  
 10-benzoyl-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
 azepino[4',5':4,5]pyrrolo-[3,2,1-ij]quinolin-4-one;  
 10-benzoyl-2-fluoro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
 5 azepino[4',5':4,5]-pyrrolo[3,2,1-ij]quinolin-4-one;  
 10-benzoyl-2,3-dichloro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
 10-benzoyl-6-propyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-  
 [3,2,1-ij]quinolin-4-one;  
 10 10-benzoyl-6-(trifluoromethyl)-5,6,9,10,11,12-hexahydro-4H,8H-  
 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
 10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol; and  
 10-benzoyl-4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
 15 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline.

All cited publications, patents, and patent documents are  
 incorporated by reference herein, as though individually incorporated by  
 reference. The invention has been described with reference to various specific  
 20 and preferred embodiments and techniques. However, it should be understood  
 that many variations and modifications may be made while remaining within the  
 spirit and scope of the invention.

## CLAIMS

What is claimed is:

1. A compound of formula I:



(I)

wherein,

each  $R_1$  is independently hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl,  $-S(O)_mNR_4R_b$ ,  $NR_4R_b$ ,  $-S(O)_mR_e$ , or  $-C(=O)NR_4R_b$ , wherein any  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, or  $C_{1-7}$ alkanoyloxy of  $R_1$  is optionally partially unsaturated and is optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy,  $-S(O)_mR_e$ ,  $-S(O)_mNR_4R_b$ ,  $NR_4R_b$ , or  $-C(=O)NR_4R_b$ ;

$R_2$  is hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl, arylcarbonyl, aryl, (aryl) $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxycarbonyl, aryloxycarbonyl, arylsulfonyl, or (aryl) $C_{1-7}$ alkoxycarbonyl;

X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy ( $-O-$ ), thio ( $-S-$ ), sulfinyl ( $-SO-$ ), sulfonyl ( $S(O)_2-$ ), or  $NR_4$  in the chain; wherein the chain is optionally substituted on each carbon with oxo ( $=O$ ), thioxo ( $=S$ ),  $-NR_4R_n$ ,  $-S(O)_pR_s$ , or  $-OR_6$ , or with one or two substituents independently selected from the group consisting of  $C_{1-7}$ alkyl,  $(C_{1-7}$ alkoxy) $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, and (aryl)oxy $C_{1-7}$ alkyl; or wherein the chain is optionally substituted on a carbon

with a 4, 5, or 6 membered spirocyclic carbon ring; or wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain (e.g. -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, or -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-) forming a ring that is fused to the ring comprising X and Y;

- 5                   each m is independently 0, 1, or 2;  
                   n is 0, 1, 2, or 3;  
                   p is 0, 1, or 2;  
                   each R<sub>a</sub> and R<sub>b</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, or (heteroaryl)C<sub>1-7</sub>alkyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;
- 10                   each R<sub>c</sub> and R<sub>d</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, C<sub>1-7</sub>alkanoyl, C<sub>1-7</sub>alkoxycarbonyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, arylcarbonyl, heteroarylcarbonyl, aryloxycarbonyl, or heteroaryloxycarbonyl; or
- 15                   R<sub>c</sub> and R<sub>d</sub> together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;
- each R<sub>e</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, or (heteroaryl)C<sub>1-7</sub>alkyl;
- R<sub>f</sub> is hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, or is a bond to a 2, 3, or 4 membered alkylene chain that
- 20                   forms a ring that is fused to the ring comprising X and Y;
- each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, or (heteroaryl)C<sub>1-7</sub>alkyl; or R<sub>g</sub> and R<sub>h</sub> together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino,
- 25                   or thiomorpholino ring;
- R<sub>i</sub> is C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, or (heteroaryl)C<sub>1-7</sub>alkyl; and
- R<sub>j</sub> is hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, or (heteroaryl)C<sub>1-7</sub>alkyl;

- wherein any aryl or heteroaryl ring of  $R_1$ ,  $R_2$ ,  $X$ ,  $Y$ ,  $R_a-R_b$ , or  $R_q-R_t$  is optionally substituted with one or more (e.g. 1, 2, 3, or 4) substituents independently selected from halo, hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy, phenyl, sulfonyl,  $NR_jR_k$ , or -
- 5  $C(=O)NR_jR_k$ ; wherein each  $R_j$  and  $R_k$  is independently hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, aryl, (aryl) $C_{1-7}$ alkyl, arylcarbonyl, or aryloxycarbonyl; or  $R_j$  and  $R_k$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring; or a pharmaceutically acceptable salt thereof;
- 10 provided  $Y$  is not oxy, thio, sulfinyl, or  $NR_p$ ; and provided  $X$  and  $Y$  together are not a 2-membered unsaturated chain; and provided no carbon of  $X$  and  $Y$  is bonded to more than one oxy, thio, sulfinyl, or  $NR_p$ .
- 15
2. The compound of claim 1 wherein each  $R_1$  is independently hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl, - $S(O)_mNR_aR_b$ ,  $NR_cR_d$ , - $S(O)_mR_e$ , or - $C(=O)NR_aR_b$ , wherein any  $C_{1-7}$ alkyl,
- 20  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, or  $C_{1-7}$ alkanoyloxy of  $R_1$  is optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, - $S(O)_mR_e$ , - $S(O)_mNR_aR_b$ ,  $NR_cR_d$ , or - $C(=O)NR_aR_b$ .
- 25
3. The compound of claim 1 wherein each  $R_1$  is independently hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl, - $S(O)_mNR_aR_b$ ,  $NR_cR_d$ , - $S(O)_mR_e$ , or - $C(=O)NR_aR_b$ , wherein any  $C_{1-7}$ alkyl or  $C_{1-7}$ alkoxy of  $R_1$  is optionally substituted with aryl, aryloxy, heteroaryl,
- 30 heteroaryloxy, hydroxy, nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,

$C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy,  $-S(O)_mR_e$ ,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ , or  $-C(=O)NR_aR_b$ .

4. The compound of claim 1 wherein each  $R_1$  is independently  
 5 hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  
 $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ ,  $-S(O)_mR_e$ , or  $-C(=O)NR_aR_b$ , wherein any  $C_{1-7}$ alkyl is  
 optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy,  
 nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy,  
 10  $-S(O)_mR_e$ ,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ , or  $-C(=O)NR_aR_b$ .

5. The compound of claim 1 wherein each  $R_1$  is independently  
 hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  
 $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ ,  $-S(O)_mR_e$ , or  $-C(=O)NR_aR_b$ .  
 15

6. The compound of claim 1 wherein each  $R_1$  is independently  
 $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy, trifluoromethyl, or halo.

- 20 7. The compound of claim 1 wherein  $n$  is 1 and  $R_1$  is  $C_{1-7}$ alkyl,  
 $C_{1-7}$ alkoxy, or halo.

8. The compound of claim 1 wherein  $n$  is 1 and  $R_1$  is methyl,  
 methoxy, chloro, or fluoro.  
 25

9. The compound of claim 1 wherein  $R_2$  is hydrogen.

10. The compound of claim 1 wherein  $n$  is 1, 2, or 3.

- 30 11. The compound of claim 1 wherein  $n$  is 0.

12. The compound of claim 1 wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (-S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), thioxo (=S), -NR<sub>q</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, or -OR<sub>p</sub>, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.
13. The compound of claim 1 wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), thioxo (=S), -NR<sub>q</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, or -OR<sub>p</sub>.
14. The compound of claim 1 wherein X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy (-O-), thio (-S-), sulfinyl (-SO-), sulfonyl (S(O)<sub>2</sub>-), or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.
15. The compound of claim 1 wherein X and Y together are a 2, 3, or 4 membered chain comprising one or more carbon atoms and optionally comprising one oxy, thio, sulfinyl, sulfonyl, or NR<sub>r</sub> in the chain; wherein the chain is optionally substituted on each carbon with oxo (=O), hydroxy, (aryl)oxy, heteroaryl(oxy) or C<sub>1-7</sub>alkoxy, or with one or two substituents independently

selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl; and wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain forming a ring that is fused to the ring comprising X and Y.

16. The compound of claim 1 wherein X and Y together are a 2, 3, or 4 membered carbon chain wherein the chain is optionally substituted on each carbon with oxo, hydroxy, (aryl)oxy, heteroaryl(oxy) or C<sub>1-7</sub>alkoxy, or with one or two substituents independently selected from the group consisting of C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.

17. The compound of claim 1 wherein X and Y together are a 2 or 3 membered carbon chain optionally substituted on each carbon with oxo or hydroxy, or with one or two C<sub>1-7</sub>alkyl.

18. The compound of claim 1 wherein X is -O-, -S-, or -C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein R<sub>g</sub> and R<sub>h</sub> are each independently hydrogen, C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl or (aryl)oxyC<sub>1-7</sub>alkyl, or R<sub>g</sub> and R<sub>h</sub> together are oxo.

19. The compound of claim 1 wherein Y is -C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)-, or -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, and each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

20. The compound of claim 1 wherein X is -O-, -S-, or -C(R<sub>g</sub>)(R<sub>h</sub>)-; and Y is -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, or -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

21. The compound of claim 1 wherein X is -O- or -S-; and Y is -C(R<sub>g</sub>)(R<sub>h</sub>)C(=O)-, -C(=O)C(R<sub>g</sub>)(R<sub>h</sub>)-, or -C(R<sub>g</sub>)(R<sub>h</sub>)C(R<sub>g</sub>)(R<sub>h</sub>)-, wherein each R<sub>g</sub> and R<sub>h</sub> is independently hydrogen or C<sub>1-7</sub>alkyl.

5

22. The compound of claim 1 wherein X and Y together are -

-CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>-,  
 -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH<sub>2</sub>-, -CH=CHCH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH=CHCH<sub>2</sub>-,  
 -O-CH<sub>2</sub>CH<sub>2</sub>-, -O-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -S-CH<sub>2</sub>CH<sub>2</sub>-, -S-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-,  
 10 -S(O)-CH<sub>2</sub>CH<sub>2</sub>-, -S(O)-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -S(O)<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>-,  
 -S(O)<sub>2</sub>-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -NR<sub>r</sub>-CH<sub>2</sub>CH<sub>2</sub>-, -NR<sub>r</sub>-CH<sub>2</sub>CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>C(=O)-,  
 -CH(R<sub>g</sub>)CH<sub>2</sub>C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>C(=O)-, -CH<sub>2</sub>OC(=O)-, -  
 CH(R<sub>g</sub>)CH<sub>2</sub>OC(=O)-, -OCH<sub>2</sub>C(=O)-, or -OCH<sub>2</sub>CH<sub>2</sub>C(=O)-; wherein each R<sub>g</sub> is  
 independently -NR<sub>r</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, or -OR<sub>r</sub>.

15

23. The compound of claim 1 wherein X and Y together are -

-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -CH=CHCH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-, -  
 -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 20 -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-,  
 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)OC(=O)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)OC(=O)-,  
 25 -OCH(R<sub>g</sub>)C(=O)-, or -OCH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-; wherein each R<sub>g</sub> is independently  
 hydrogen, C<sub>1-7</sub>alkyl, aryl, (aryl)C<sub>1-7</sub>alkyl, (aryl)oxy, heteroaryl(oxy), or  
 (aryl)oxyC<sub>1-7</sub>alkyl.

24. The compound of claim 1 wherein X and Y together are -

30 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-,



-O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)C(=O)-,  
 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-, -CH(R<sub>g</sub>)OC(=O)-, -OCH(R<sub>g</sub>)C(=O)-; wherein each R<sub>g</sub> is  
 independently hydrogen or C<sub>1-7</sub>alkyl.

5

25. The compound of claim 1 wherein X and Y together are -  
 -CH(R<sub>g</sub>)CH<sub>2</sub>-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH<sub>2</sub>-, or -O-CH<sub>2</sub>CH<sub>2</sub>-, wherein each R<sub>g</sub> is  
 independently -NR<sub>q</sub>R<sub>r</sub>, -S(O)<sub>p</sub>R<sub>s</sub>, -OR<sub>v</sub>, C<sub>1-7</sub>alkyl, (C<sub>1-7</sub>alkoxy)C<sub>1-7</sub>alkyl, aryl,  
 (aryl)C<sub>1-7</sub>alkyl, heteroaryl, (heteroaryl)C<sub>1-7</sub>alkyl, and (aryl)oxyC<sub>1-7</sub>alkyl.

10

26. The compound of claim 1 wherein X and Y together are -  
 -C(=O)CH<sub>2</sub>-, -CH<sub>2</sub>C(=O)-, -C(=S)CH<sub>2</sub>-, -CH<sub>2</sub>C(=S)-, -C(=O)CH<sub>2</sub>CH<sub>2</sub>-,  
 -CH<sub>2</sub>C(=O)CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>C(=O)-, -C(=S)CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>C(=S)CH<sub>2</sub>-, or  
 -CH<sub>2</sub>CH<sub>2</sub>C(=S)-.

15

27. The compound of claim 1 wherein X and Y together are -  
 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -C(R<sub>g</sub>)=C(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -S(O)-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 -S(O)<sub>2</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -NR<sub>r</sub>-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, or -CH(R<sub>g</sub>)CH(R<sub>g</sub>)C(=O)-;  
 20 wherein each R<sub>g</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, or together with an R<sub>g</sub> on  
 an adjacent carbon atom forms a fused 4, 5, or 6, membered carbocyclic ring.

28. The compound of claim 1 wherein X and Y together are -  
 -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-,  
 25 -S-CH(R<sub>g</sub>)CH(R<sub>g</sub>)-; wherein each R<sub>g</sub> is independently hydrogen, C<sub>1-7</sub>alkyl, aryl,  
 or (aryl)C<sub>1-7</sub>alkyl.

29. The compound of claim 1 wherein X and Y together are -  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>C(CH<sub>3</sub>)H-, -CH<sub>2</sub>C(CH<sub>3</sub>)HCH<sub>2</sub>-, -C(CH<sub>3</sub>)HCH<sub>2</sub>CH<sub>2</sub>-, -

-CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>C(CH<sub>3</sub>)H-, -C(CH<sub>3</sub>)HC(CH<sub>3</sub>)H-, -CH(R<sub>g</sub>)CH(R<sub>g</sub>)-, -O-CH<sub>2</sub>CH<sub>2</sub>-,  
 -O-C(CH<sub>3</sub>)HCH<sub>2</sub>-,  
 or -S-CH<sub>2</sub>CH<sub>2</sub>-.

5 30. The compound of claim 1 wherein X and Y together are -

-CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH=CHCH<sub>2</sub>-, -O-CH<sub>2</sub>CH<sub>2</sub>-, -S-CH<sub>2</sub>CH<sub>2</sub>-,  
 -S(O)-CH<sub>2</sub>CH<sub>2</sub>-, -S(O)<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>-, -NR<sub>f</sub>-CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>C(=O)-,  
 -CH<sub>2</sub>CH<sub>2</sub>C(=O)-, -CH<sub>2</sub>OC(=O)-, -OCH<sub>2</sub>C(=O)-.

10 31. The compound of any one of claims 1 to 30 wherein R<sub>2</sub> is  
 hydrogen

32. The compound of claim 1 which is:

5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

15 ij]quinoline;

2-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

ij]quinoline;

1-methoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

ij]quinoline;

20 2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

ij]quinoline;

6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

ij]quinoline;

5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

25 ij]quinoline;

4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-

ij]quinoline;

(+)-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-

azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;

- (-)-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- (+)-5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5 (-)-5-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- (+)-4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- (-)-4-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 15 5-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 4,5-dimethyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2,3,4,5,8b,9,10,11,12,12a-decahydro-1H-azepino[4',5':4,5]pyrrolo[3,2,1-jk]carbazole;
- 20 6-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 6-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 25 5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 5-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 30 hi]indole;

- 2-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]thiazino[2,3,4-hi]indole;
- 2,2-dimethyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 5 4-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 4-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 10 5-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 6-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 5,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 15 5-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 6-chloro-5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 20 5,6-dichloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 4,6-dichloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole;
- 1-chloro-2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 25 1-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;

- 2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 2-chloro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 5 6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 2-fluoro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 2,3-dichloro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 10 6-propyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 6-(trifluoromethyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;
- 15 5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol;
- 4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10-benzoyl-4-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline; or
- 20 4-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- or a pharmaceutically acceptable salt thereof.
- 25 33. The compound 1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole; or a pharmaceutically acceptable salt thereof.
34. The compound of claim 1 which is:
- 4-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4]oxazino[2,3,4-
- 30 hi]indole;

- 5-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 6-bromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 5 5,6-dibromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 4,6-dibromo-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 4-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 10 5-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 6-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 15 4-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 5-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 4-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 20 5-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 6-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] oxazino[2,3,4-hi]indole;
- 25 4-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 30

- 4-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5 6-chloro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 4,5-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 10 4,6-difluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 4-chloro-5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 15 4-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5-chloro-6-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 6-chloro-5-fluoro-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 20 4-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 25 6-methyl-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 4-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 30 hi]indole;

- 6-methoxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 4-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 5 5-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 6-(trifluoromethyl)-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 4-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 10 5-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 6-benzyloxy-1,2,8,9,10,11-hexahydro-7H-azepino[4,5-b][1,4] thiazino[2,3,4-hi]indole;
- 15 1-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 3-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 1-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 3-bromo-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 20 3-chloro-1-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-chloro-1-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 1-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 3-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 25 4-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 1-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 2-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;



- 3-(trifluoromethyl)-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;
- 4-benzyl-oxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5 4-(3-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(2-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(4-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 4-(3-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 15 4-(4-bromo-2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- N-phenyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- 20 N-(4-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- N-(3-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- 25 N-(2-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- N-(4-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- N-(3-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- 30

- N-(2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- N-(4-bromo-2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-amine;
- 5 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-4-thione;
- 4-(phenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(4-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 4-(3-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(2-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 15 4-(4-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(3-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 4-(2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 20 4-(4-bromo-2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-ol;
- 25 5-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-benzyloxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-phenoxy-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 30 ij]quinoline;

- 5-(4-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-(3-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5 5-(2-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-(4-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-(3-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 5-(2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5-(4-bromo-2-methoxyphenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 15 5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- N-phenyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- N-(4-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- 20 N-(3-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- N-(2-chlorophenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- 25 N-(4-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- N-(3-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- N-(2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;
- 30

- N-(4-bromo-2-methoxyphenyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-5-amine;  
5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-5-thione;
- 5 5-(phenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5-(4-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5-(3-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 5-(2-chlorophenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5-(4-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 15 5-(3-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5-(2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5-(4-bromo-2-methoxyphenylsulfonyl)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 20 4-(4-chlorophenoxy)-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-[2-(4-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 25 1-[2-(3-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-[2-(2-fluorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-[2-(4-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 30

- 1-[2-(3-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(2-chlorophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 5 1-[2-(4-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(3-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(2-bromophenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 10 1-[2-(4-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(3-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 15 1-[2-(2-methoxyphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(4-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(3-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 20 1-[2-(2-methylphenoxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-(1-naphthyloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 25 1-[2-(2-naphthyloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-([1,1'-biphenyl]-4-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 1-[2-([1,1'-biphenyl]-3-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;
- 30

- 1-[2-([1,1'-biphenyl]-2-yloxy)ethoxy]-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-{2-[4-(trifluoromethoxy)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
5 1-{2-[3-(trifluoromethoxy)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-{2-[2-(trifluoromethoxy)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
1-{2-[4-(trifluoromethyl)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline;  
10 1-{2-[3-(trifluoromethyl)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline; or  
1-{2-[2-(trifluoromethyl)phenoxy]ethoxy}-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline; or a pharmaceutically acceptable salt  
15 thereof.

35. A pharmaceutical composition comprising a compound of any one of claims 31 to 34 and a pharmaceutically acceptable excipient.
- 20 36. A compound of any one of claims 31 to 34 for use in medical diagnosis or therapy.
37. The compound of claim 36 wherein the therapy is the treatment of a disease or disorder of the central nervous system.
- 25 38. The compound of claim 37 wherein the disease or disorder of the central nervous system is obesity, depression, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, a stress related disease, panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress

syndrome, immune system depression, a stress induced problem with the urinary, gastrointestinal or cardiovascular system, neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine, headaches, cluster headaches, sexual dysfunction, addictive disorder and withdrawal  
5 syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to general medical conditions, attention-deficit hyperactivity disorder, behavioral disturbance, bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and  
10 other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, movement disorder, oppositional defiant disorder, peripheral neuropathy, post-traumatic stress disorder, premenstrual dysphoric disorder, a psychotic disorder, mood disorder, seasonal affective disorder, a sleep disorder, a specific development disorder, agitation disorder, selective serotonin  
15 reuptake inhibition syndrome, or a Tic disorder.

39. The compound of claim 36 wherein the therapy is the treatment of anxiety, obesity, depression, or a stress related disease.

20 40. The use of a compound of any one of claims 31 to 34 to prepare a medicament for treating or preventing a disease or disorder of the central nervous system.

41. The use of claim 40 wherein the disease or disorder of the central  
25 nervous system is obesity, depression, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, a stress related disease, panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, a stress induced problem with the urinary, gastrointestinal or cardiovascular system, neurodegenerative disorders,  
30 autism, chemotherapy-induced vomiting, hypertension, migraine, headaches,

cluster headaches, sexual dysfunction, addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to general medical conditions, attention-deficit hyperactivity disorder, behavioral  
5 disturbance, bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, movement disorder, oppositional defiant disorder, peripheral neuropathy, post-traumatic stress disorder, premenstrual dysphoric  
10 disorder, a psychotic disorder, mood disorder, seasonal affective disorder, a sleep disorder, a specific development disorder, agitation disorder, selective serotonin reuptake inhibition syndrome, or a Tic disorder.

42. The use of claim 40 wherein the disease or disorder of the central  
15 nervous system is anxiety, obesity, depression, or a stress related disease.

43. A method for treating a disease or condition in a mammal wherein the 5-HT receptor is implicated and modulation of 5-HT function is desired comprising administering a therapeutically effective amount of a  
20 compound of any one of claims 31 to 35 to the mammal.

44. The method of claim 43 wherein the disease or condition is anxiety, obesity, depression, schizophrenia, a stress related disease, panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress  
25 syndrome, immune system depression, a stress induced problem with the gastrointestinal or cardiovascular system, or sexual dysfunction.


45. The method of claim 44 wherein the disease is anxiety, obesity, depression, or a stress related disease.



46. A method for treating or preventing a disease or disorder of the central nervous system in a mammal comprising administering a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof to the mammal.

- 5 47. The method of claim 46 wherein the disease or disorder of the central nervous system is obesity, depression, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, a stress related disease, panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, a stress induced problem with the  
10 urinary, gastrointestinal or cardiovascular system, neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine, headaches, cluster headaches, sexual dysfunction, addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to general  
15 medical conditions, attention-deficit hyperactivity disorder, behavioral disturbance, bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, movement disorder, oppositional defiant disorder,  
20 peripheral neuropathy, post-traumatic stress disorder, premenstrual dysphoric disorder, a psychotic disorder, mood disorder, seasonal affective disorder, a sleep disorder, a specific development disorder, agitation disorder, selective serotonin reuptake inhibition syndrome, or a Tic disorder.

- 25 48. A compound of formula I:



(I)

wherein,

- each  $R_1$  is independently hydroxy, nitro, halo, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy, aryl, heteroaryl,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ ,  $-S(O)_mR_e$ , or  $-C(=O)NR_aR_b$ , wherein any  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, or  $C_{1-7}$ alkanoyloxy of  $R_1$  is optionally partially unsaturated and is optionally substituted with aryl, aryloxy, heteroaryl, heteroaryloxy, hydroxy, nitro, halo, cyano,  $C_{1-7}$ alkoxy,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl,  $C_{1-7}$ alkanoyloxy,  $-S(O)_mR_e$ ,  $-S(O)_mNR_aR_b$ ,  $NR_cR_d$ , or  $-C(=O)NR_aR_b$ ;
- $R_2$  is a suitable protecting group;
- X and Y together are a 2, 3, or 4 membered saturated or partially unsaturated chain comprising one or more carbon atoms and optionally comprising one oxy ( $-O-$ ), thio ( $-S-$ ), sulfinyl ( $-SO-$ ), sulfonyl ( $S(O)_2-$ ), or  $NR_f$  in the chain; wherein the chain is optionally substituted on each carbon with oxo ( $=O$ ), thioxo ( $=S$ ),  $-NR_gR_h$ ,  $-S(O)_pR_s$ , or  $-OR_t$ , or with one or two substituents independently selected from the group consisting of  $C_{1-7}$ alkyl,  $(C_{1-7}alkoxy)C_{1-7}alkyl$ , aryl,  $(aryl)C_{1-7}alkyl$ , heteroaryl,  $(heteroaryl)C_{1-7}alkyl$ , and  $(aryl)oxyC_{1-7}alkyl$ ; or wherein the chain is optionally substituted on a carbon with a 4, 5, or 6 membered spirocyclic carbon ring; or wherein the chain is optionally substituted on two adjacent atoms with a 2, 3, or 4 membered alkylene chain (e.g.  $-CH_2CH_2-$ ,  $-CH_2CH_2CH_2-$ , or  $-CH_2CH_2CH_2CH_2-$ ) forming a ring that is fused to the ring comprising X and Y;
- each m is independently 0, 1, or 2;
- n is 0, 1, 2, or 3;
- p is 0, 1, or 2;
- each  $R_a$  and  $R_b$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl,  $(aryl)C_{1-7}alkyl$ , heteroaryl, or  $(heteroaryl)C_{1-7}alkyl$ ; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

each  $R_c$  and  $R_d$  is independently hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, arylcarbonyl, heteroarylcarbonyl, aryloxy carbonyl, or heteroaryloxy carbonyl; or  $R_c$  and  $R_d$  together with the nitrogen to which they are attached form a

5 pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

each  $R_e$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl;

$R_f$  is hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, (heteroaryl) $C_{1-7}$ alkyl, or is a bond to a 2, 3, or 4 membered alkylene chain that

10 forms a ring that is fused to the ring comprising X and Y;

each  $R_g$  and  $R_h$  is independently hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl; or  $R_g$  and  $R_h$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

15  $R_i$  is  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl; and

$R_j$  is hydrogen,  $C_{1-7}$ alkyl, aryl, (aryl) $C_{1-7}$ alkyl, heteroaryl, or (heteroaryl) $C_{1-7}$ alkyl;

wherein any aryl or heteroaryl ring of  $R_1$ ,  $R_2$ , X, Y,  $R_4$ - $R_n$ , or  $R_q$ - $R_t$

20 is optionally substituted with one or more (e.g. 1, 2, 3, or 4) substituents independently selected from halo, hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkoxy, phenyl, sulfonyl,  $NR_jR_k$ , or - $C(=O)NR_jR_k$ ; wherein each  $R_j$  and  $R_k$  is independently hydrogen,  $C_{1-7}$ alkyl,  $C_{1-7}$ alkanoyl,  $C_{1-7}$ alkoxycarbonyl, aryl, (aryl) $C_{1-7}$ alkyl, arylcarbonyl, or

25 aryloxy carbonyl; or  $R_j$  and  $R_k$  together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, or thiomorpholino ring;

or a pharmaceutically acceptable salt thereof;

provided Y is not oxy, thio, sulfinyl, or  $NR_p$ ; and

provided X and Y together are not a 2-membered unsaturated

30 chain; and

provided no carbon of X and Y is bonded to more than one oxy,  
thio, sulfinyl, or NR<sub>r</sub>.

49. The compound of claim 1 wherein R<sub>2</sub> is C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl,  
5 arylcarbonyl, (aryl)C<sub>1-2</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, aryloxycarbonyl, arylsulfonyl,  
or (aryl)methoxycarbonyl, wherein any aryl is optionally substituted with 1, 2, or  
3 substituents independently selected from C<sub>1-4</sub>alkyl and trifluoromethyl.
50. The compound of claim 1 wherein R<sub>2</sub> is methyl, ethyl, propyl,  
10 isopropyl, acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, benzyl, or *p*-  
toluenesulfonyl.
51. The compound of claim 1 which is: 10-benzoyl-5,6,9,10,11,12-  
hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
15 10-benzoyl-2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-2-chloro-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
20 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-2-fluoro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-2,3-dichloro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
25 10-benzoyl-6-propyl-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-6-(trifluoromethyl)-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
30 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol; or

10-benzoyl-4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline.

52. The compound of claim 1 which is: benzyl 5,6,8,9,11,12-  
5 hexahydro-4H,10H-azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
benzyl 2-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
10-benzoyl-1-methoxy-5,6,9,10,11,12-hexahydro-4H,8H-  
azepino[4',5':4,5]pyrrolo-[3,2,1-ij]quinoline;  
10 benzyl 2-fluoro-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
15 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(+)-benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
20 (-)-benzyl 6-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(+)-benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(-)-benzyl 5-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
25 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(+)-benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;  
(-)-benzyl 4-methyl-5,6,8,9,11,12-hexahydro-4H,10H-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline-10-carboxylate;

- benzyl 2-methyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 2,2-dimethyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 5 benzyl 4-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 4-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 5-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 10 benzyl 5-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 5-methyl-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 15 benzyl 6-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 6-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 6-methyl-1,2,6b,7,8,10,11,11a-octahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 20 benzyl 6-(trifluoromethyl)-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 5,6-difluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylatebenzyl;
- 25 benzyl 5-chloro-6-fluoro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 5-fluoro-6-chloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- benzyl 5,6-dichloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;
- 30

- benzyl 4,6-dichloro-1,2,7,8,10,11-hexahydro-9H-azepino[4,5-b][1,4]oxazino[2,3,4-hi]indole-9-carboxylate;  
 7-benzoyl 4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]indole;  
 7-benzoyl 2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
 5 hi]indole;  
 7-benzoyl 2-methoxy-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
 hi]indole;  
 benzyl-5-methyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-hi]-7-  
 carboxylate;  
 10 benzyl-4,5-dimethyl-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
 hi]indole-7-carboxylate;  
 benzyl-2,3,4,5,8b,9,10,11,12,12a-decahydro-1H-azepino[4',5':4,5]pyrrolo[3,2,1-  
 jk]carbazole-7-carboxylate;  
 benzyl-1-chloro-2-fluoro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-  
 15 b]pyrrolo[3,2,1-hi]indole-7-carboxylate;  
 benzyl-2-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
 hi]indole-7-carboxylate;  
 benzyl-1-chloro-4,5,8,9,10,11-hexahydro-7H-azepino[4,5-b]pyrrolo[3,2,1-  
 hi]indole-7-carboxylate;  
 20 10-benzoyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo[3,2,1-  
 ij]quinolin-4-one;  
 10-benzoyl-2-fluoro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-  
 [3,2,1-ij]quinolin-4-one;  
 10-benzoyl-2-chloro-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-  
 25 [3,2,1-ij]quinolin-4-one;  
 10-benzoyl-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
 azepino[4',5':4,5]pyrrolo-[3,2,1-ij]quinolin-4-one;  
 10-benzoyl-2-fluoro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-  
 azepino[4',5':4,5]-pyrrolo[3,2,1-ij]quinolin-4-one;  
 30 10-benzoyl-2,3-dichloro-6-methyl-5,6,9,10,11,12-hexahydro-4H,8H-

azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-6-propyl-5,6,9,10,11,12-hexahydro-4H,8H-azepino[4',5':4,5]pyrrolo-  
[3,2,1-ij]quinolin-4-one;  
10-benzoyl-6-(trifluoromethyl)-5,6,9,10,11,12-hexahydro-4H,8H-  
5 azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-one;  
10-benzoyl-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinolin-4-ol; and  
10-benzoyl-4-methoxy-5,6,8,9,10,11,12,12a-octahydro-4H,7aH-  
azepino[4',5':4,5]pyrrolo[3,2,1-ij]quinoline.

10

53. A method for preparing a compound of formula (I) wherein R<sub>2</sub> is hydrogen comprising deprotecting a corresponding compound of formula (I) wherein R<sub>2</sub> is a protecting group.

15 54. The method of claim 53 wherein the protecting group is benzyloxycarbonyl, or benzoyl.








Figure 1





Figure 2

Figure 3




Figure 4




Figure 5





Figure 6



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/10600

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D471/16 A61K31/395 A61P43/00 C07D498/16 C07D513/16  
/(C07D471/16,223:00,221:00,209:00),(C07D471/16,223:00,209:00,  
209:00),(C07D498/16,265:00,223:00,209:00),(C07D513/16,279:00,

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 377 238 A (DUPHAR INTERNATIONAL RESEARCH B.V.) 11 July 1990 (1990-07-11) page 3, line 22 - line 33; claim 1	1,35-37
A	US 3 839 357 A (JACKSON B. HESTER ET AL.) 1 October 1974 (1974-10-01) cited in the application column 1 -column 2	1,35,36
A	US 3 676 558 A (JACKSON B. HESTER ) 11 July 1972 (1972-07-11) cited in the application column 1 -column 2	1,35,36

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 September 2000

Date of mailing of the international search report

04/10/2000

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10600

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 223:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*8\* document member of the same patent family

Date of the actual completion of the international search

12 September 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10600

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 377238 A	11-07-1990	AU 615818 B	10-10-1991
		AU 4716589 A	28-06-1990
		CA 2005974 A	22-06-1990
		DK 647389 A	23-06-1990
		JP 2258785 A	19-10-1990
		NZ 231870 A	25-02-1992
		US 5223625 A	29-06-1993
		ZA 8909744 A	26-09-1990
US 3839357 A	01-10-1974	BE 698268 A	10-11-1967
		CH 506543 A	30-04-1971
		DE 1695943 A	13-05-1971
		ES 340258 A	01-06-1968
		FR 6699 M	10-02-1969
		FR 1524495 A	02-09-1968
		GB 1180615 A	04-02-1970
		GB 1180616 A	04-02-1970
		IL 27776 A	29-11-1971
		IL 36801 A	29-11-1971
		NL 6706513 A	13-11-1967
US 3676558 A	11-07-1972	NONE	

**THIS PAGE BLANK (USPTO)**